

Quantum Mechanics: Commutation Relation Proofs

5th April 2010

I. Proof for Non-Commutativity of Individual Quantum Angular Momentum Operators

In this section, we will show that the operators \hat{L}_x , \hat{L}_y , \hat{L}_z do not commute with one another, and hence cannot be known simultaneously. The relations are (reiterating from previous lectures):

$$\hat{L}_x = -i\hbar \left(y \frac{\partial}{\partial z} - z \frac{\partial}{\partial y} \right)$$

$$\hat{L}_y = -i\hbar \left(z \frac{\partial}{\partial x} - x \frac{\partial}{\partial z} \right)$$

$$\hat{L}_z = -i\hbar \left(x \frac{\partial}{\partial y} - y \frac{\partial}{\partial x} \right)$$

We would like to prove the following commutation relations:

$$[\hat{L}_x, \hat{L}_y] = i\hbar \hat{L}_z,$$

$$[\hat{L}_y, \hat{L}_z] = i\hbar \hat{L}_x,$$

$$[\hat{L}_z, \hat{L}_x] = i\hbar \hat{L}_y.$$

We will use the first relation for our proof; the second and third follow analogously. Let's also consider a function, $f(x, y, z)$ that we will have the operators act upon in our discussion. The expanded version of $[\hat{L}_x, \hat{L}_y] = i\hbar \hat{L}_z$ is:

$$[\hat{L}_x, \hat{L}_y] = \left((\hat{L}_x) (\hat{L}_y) - (\hat{L}_y) (\hat{L}_x) \right)$$

$$= \left(\left(y \frac{\partial}{\partial z} - z \frac{\partial}{\partial y} \right) \left(z \frac{\partial}{\partial x} - x \frac{\partial}{\partial z} \right) \right) - \left(\left(z \frac{\partial}{\partial x} - x \frac{\partial}{\partial z} \right) \left(y \frac{\partial}{\partial z} - z \frac{\partial}{\partial y} \right) \right)$$

Now let's expand the operators (ignore the underbraces and numbers for now):

$$\begin{aligned} (\hat{L}_x)(\hat{L}_y) &= \left(y \frac{\partial}{\partial z} - z \frac{\partial}{\partial y} \right) \left(z \frac{\partial}{\partial x} - x \frac{\partial}{\partial z} \right) \\ &= y \frac{\partial}{\partial z} \left(z \frac{\partial}{\partial x} \right) - y \frac{\partial}{\partial z} \left(x \frac{\partial}{\partial z} \right) - z \frac{\partial}{\partial y} \left(z \frac{\partial}{\partial x} \right) + z \frac{\partial}{\partial y} \left(x \frac{\partial}{\partial z} \right) \\ &= \underbrace{y \left(\frac{\partial}{\partial x} + z \frac{\partial}{\partial z} \frac{\partial}{\partial x} \right)}_{\mathbf{1}} - \underbrace{y \left(x \frac{\partial}{\partial z} \frac{\partial}{\partial z} \right)}_{\mathbf{2}} - \underbrace{z \left(z \frac{\partial}{\partial y} \frac{\partial}{\partial x} \right)}_{\mathbf{3}} + \underbrace{z \left(x \frac{\partial}{\partial y} \frac{\partial}{\partial z} \right)}_{\mathbf{4}} \end{aligned}$$

Likewise for the $(\hat{L}_y)(\hat{L}_x)$ term:

$$\begin{aligned} (\hat{L}_y)(\hat{L}_x) &= \left(z \frac{\partial}{\partial x} - x \frac{\partial}{\partial z} \right) \left(y \frac{\partial}{\partial z} - z \frac{\partial}{\partial y} \right) \\ &= z \frac{\partial}{\partial x} \left(y \frac{\partial}{\partial z} \right) - z \frac{\partial}{\partial x} \left(z \frac{\partial}{\partial y} \right) - x \frac{\partial}{\partial z} \left(y \frac{\partial}{\partial z} \right) + x \frac{\partial}{\partial z} \left(z \frac{\partial}{\partial y} \right) \\ &= \underbrace{z \left(y \frac{\partial}{\partial x} \frac{\partial}{\partial z} \right)}_{\mathbf{5}} - \underbrace{z \left(z \frac{\partial}{\partial x} \frac{\partial}{\partial y} \right)}_{\mathbf{6}} - \underbrace{x \left(y \frac{\partial}{\partial z} \frac{\partial}{\partial z} \right)}_{\mathbf{7}} + \underbrace{x \left(\frac{\partial}{\partial y} + z \frac{\partial}{\partial z} \frac{\partial}{\partial y} \right)}_{\mathbf{8}} \end{aligned}$$

Combining terms and noting that $[\hat{x}, \hat{y}] = 0$ and $[\hat{p}_x, \hat{p}_y] = 0$. We will combine terms as follows: (1-8), (2-7), (3-6), and (4-5).

$$\begin{aligned} (1-8) &= y \left(\frac{\partial}{\partial x} + z \frac{\partial}{\partial z} \frac{\partial}{\partial x} \right) - x \left(\frac{\partial}{\partial y} + z \frac{\partial}{\partial z} \frac{\partial}{\partial y} \right) \\ &= \left(y \frac{\partial}{\partial x} - x \frac{\partial}{\partial y} \right) + \left(yz \frac{\partial}{\partial z} \frac{\partial}{\partial x} - xz \frac{\partial}{\partial z} \frac{\partial}{\partial y} \right) \\ &= \left(y \frac{\partial}{\partial x} - x \frac{\partial}{\partial y} \right) + \left(zy \frac{\partial}{\partial x} \frac{\partial}{\partial z} - zx \frac{\partial}{\partial y} \frac{\partial}{\partial z} \right) \quad (\text{recall the commutation relations noted!}) \\ &= \underbrace{\left(y \frac{\partial}{\partial x} - x \frac{\partial}{\partial y} \right) + z \left(y \frac{\partial}{\partial x} - x \frac{\partial}{\partial y} \right) \frac{\partial}{\partial z}}_{\mathbf{9}} \end{aligned}$$

$$\begin{aligned}
(2-7) &= xy \frac{\partial^2}{\partial z^2} - yx \frac{\partial^2}{\partial z^2} \\
&= (xy - yx) \frac{\partial^2}{\partial z^2} \\
&= [x, y] \frac{\partial^2}{\partial z^2} \\
&= (0) \frac{\partial^2}{\partial z^2} = 0
\end{aligned}$$

$$\begin{aligned}
(3-6) &= z^2 \frac{\partial}{\partial x} \frac{\partial}{\partial y} - z^2 \frac{\partial}{\partial y} \frac{\partial}{\partial x} \\
&= z^2 \left(\frac{\partial}{\partial x} \frac{\partial}{\partial y} - \frac{\partial}{\partial y} \frac{\partial}{\partial x} \right) \\
&= z^2 (\hat{p}_x \hat{p}_y - \hat{p}_y \hat{p}_x) \\
&= z^2 [\hat{p}_x, \hat{p}_y] = 0 \quad (\hat{p}_x \text{ and } \hat{p}_y \text{ commute!})
\end{aligned}$$

$$\begin{aligned}
(4-5) &= zx \frac{\partial}{\partial y} \frac{\partial}{\partial z} - zy \frac{\partial}{\partial x} \frac{\partial}{\partial z} \\
&= z \underbrace{\left(x \frac{\partial}{\partial y} - y \frac{\partial}{\partial x} \right)}_{\mathbf{10}} \frac{\partial}{\partial z}
\end{aligned}$$

If we add the expressions 9 and 10, inserting a factor of $-i\hbar$ for each partial derivative that represents a momentum operator, we obtain:

$$\begin{aligned}
[\hat{L}_x, \hat{L}_y] &= -\hbar^2 \left(y \frac{\partial}{\partial x} - x \frac{\partial}{\partial y} \right) \\
&= \hbar^2 \left(x \frac{\partial}{\partial y} - y \frac{\partial}{\partial x} \right)
\end{aligned}$$

But,

$$\hat{L}_z = -i\hbar \left(x \frac{\partial}{\partial y} - y \frac{\partial}{\partial x} \right)$$

Thus,

$$[\hat{L}_x, \hat{L}_y] = i\hbar \hat{L}_z$$