



Determining the Presence of Glyphosate-Resistant Horseweed Under Field Conditions

Mark VanGessel, Professor/Extension Specialist,

Weed Science and Crop Management Department of Plant and Soil Sciences, University of Delaware

October 2025 (revised)

Glyphosate-resistant horseweed (or marestalk or *Conyza canadensis*) has been confirmed throughout Delaware, Maryland and New Jersey. Horseweed is a winter annual weed. Emergence begins in the fall and continues into the early summer. The plant initially grows low to the ground (rosette stage) and then develops a tall, erect stem (bolting stage), from which numerous seed heads are produced. Resistant-horseweed can tolerate glyphosate treatments at all stages, even while the plant is in the rosette stage.

Glyphosate is the active ingredient in Roundup and numerous other brand names. If resistance is suspected in one of your fields, some questions should be asked to help determine if herbicide resistance is the reason for the lack of control.

- Was the herbicide applied as recommended?
- Did the plants emerge after application?
- Are there patterns to where weeds were not controlled in the field? (This implies a problem with the application.)

Resistance develops by selecting resistant biotypes through repeated use of the same herbicide mode of action. Some questions regarding herbicide use need to be considered:

- Has the same herbicide (or herbicides) with the same mode of action been used consistently over a period of years?
- Has the species been controlled by the herbicide(s) in the recent past?
- Has there been a decline in control of the suspected species?

One of the classic signs for resistance with postemergence herbicides is that only one weed species is present, all others are effectively controlled

(or uniformly suppressed, depending on the herbicide and weed).




Figure 1: The photo on the left is of a plant sprayed with glyphosate, showing no difference in growth due to the treatment. The photo on the right is a horseweed plant treated with glyphosate, and the top portion of the plant was killed, but the plant does not die and begins to produce long branches from lateral buds on the lower portion of the stem. Both plants are resistant to glyphosate.

The level of resistance in the horseweed populations is not as great as seen with other herbicide-resistant weeds. It is common to see some glyphosate response in the resistant horseweed. The response can be either yellowing in the growing point or death of the growing point, resulting in long branches developing from lateral buds on the lower part of the plant (see figure 1). Given the situation in our region, there should be some susceptible plants (controlled by glyphosate) in the same field along with resistant plants (surviving plants). This occurrence of controlled and uncontrolled plants randomly scattered throughout the field is an easy way to identify fields with resistant horseweed (see figure 2).



Figure 2: Randomly scattered plants throughout the field are an easy way to identify fields with resistant horseweed.

If you suspect resistance, but are still uncertain, or your situation is not covered by the scenarios described above, there is a test you can do yourself. Mix up a small hand sprayer with 1% solution of glyphosate (2.5 tablespoons per 1 gallon of water or one-half cup per 3 gallons of water) and spray the suspected plants. Spray approximately 30 plants in a small area. Do not “douse” the plants with the spray mixture; rather, a light spray is adequate. Then spray the rest of the field with the appropriate herbicide to prevent the horseweed from producing seed. This procedure will help you determine if glyphosate-resistant horseweed is present in your field, which in turn will allow better management decisions to be made. Once a field is identified with glyphosate-resistant horseweed, glyphosate will no longer control this species, and additional management will be needed for the present year and many future years.

This information is brought to you by the University of Delaware Cooperative Extension, a service of the UD College of Agriculture and Natural Resources — a land-grant institution. This institution is an equal opportunity provider.