




Hand Washing Procedures

Written by: Sue Snider, Ph.D. Professor/Food Safety and Nutrition Specialist.

Revised by: Diane Oliver, MPH, RDN, LDN

August 2025



Double-hand wash method

Use the double-hand wash method to reduce disease-causing (pathogenic) organisms on hands. Water temperature should be at least 100 F and scrubbing should be done for at least 10 seconds. The whole process should take 20 seconds.

The steps include:

1. Wet hands under hot, flowing water (100° F to 120° F, 2 gallons per minute).
2. Apply sufficient soap to develop a good lather. (Refrain from using antibacterial soaps.)
3. Use a fingernail brush to eliminate dirt on fingertips and under fingernails.
4. Rinse hands and brush in hot (100° F to 120° F), flowing (more than 2 gallons per minute) water.
5. To ensure removal of pathogens, wash hands again without using the fingernail

brush. Wash arms as far up as will make contact with food.

6. Rinse hands and arms again in hot, flowing water.
7. Dry hands and arms with disposable paper towels.

Poor personal hygiene of workers causes 25 to 40 percent of all foodborne related illness in this country. Proper attention to good personal hygiene is essential to decrease foodborne disease.

This information is brought to you by the University of Delaware Cooperative Extension, a service of the UD College of Agriculture and Natural Resources — a land-grant institution. This institution is an equal opportunity provider.