

2026/2027
Mid-Atlantic Commercial Vegetable
Production Recommendations



F. Commodity Recommendations

Pesticide Use Disclaimer

THE LABEL IS THE LAW

A pesticide applicator is legally bound by the labeling found on and with the pesticide container in their possession. Before using a pesticide, check and always follow the labeling distributed with the product at the point of sale for legally enforceable rates and use restrictions and precautions.

Although labels are available on the Internet from electronic label services such as Proagrica's CDMS (<https://www.cdms.net/>), Greenbook (<https://www.greenbook.net/>), or Agworld DBX powered by Agrian (<https://www.agrian.com/labelcenter/results.cfm>) the information contained in these electronic labels may not be identical to the labeling distributed with the product. **Please be advised that these electronic label services provide use disclaimers, and in some cases legally binding *User Agreements* assigning ALL liability to user of service.** (See section D 3.1. Labels and Labeling for more detail.)

Guide to the Recommended Pesticide Tables in the Following Crop Sections:

1. Pesticides are listed by **group number or code based on chemical structure and mechanism of action**, as classified by the Herbicide Resistance Action Committee (HRAC, <https://hracglobal.com>) for herbicides, the Insecticide Resistance Action Committee (IRAC, <https://irac-online.org>) for insecticides, and the Fungicide Resistance Action Committee (FRAC, <https://www.frac.info/>) for fungicides. **In this guide, if the group number or code is in bold font, there are resistance concerns for the product.**
2. **Restricted use pesticides** are marked with a * in the Tables. These products may only be used by certified and/or licensed pesticide applicators, and when stated on the label, those making applications under their direct supervision. Some labels may restrict use solely to certified and/or licensed applicators. (See section D 3.2.1 Restricted Use Classification Statement for more detail).
3. **In addition to the pesticide products listed in the Commodity Recommendations below, other formulations or brands with the same active ingredient(s) may be commercially available. ALWAYS CHECK THE LABELING ON THE PRODUCT CONTAINER ITSELF:**
 - a) to ensure a pesticide is labeled for the same intended use,
 - b) to ensure the pesticide is labeled for the desired crop,
 - c) for differences in application rates and % active ingredient(s), and
 - d) additional restrictions.
4. All pesticide recommendations contained in this document are prescribed for spray applications to a **broadcast area of 1 acre** (43,560 square feet). **Adjust the rate accordingly for banded applications** (See section E 1.3. Calibrating Granular Applicators) **or for chemigation** (check labels for amounts per 1,000 feet).
5. Check the physical product label for and do not exceed the maximum amount of pesticide *per application* and the maximum number of applications *per year*.
6. **Bee Toxicity Rating (Bee TR):** N=nontoxic; L=minimum impact on bees; M=moderately toxic, can be used if dosage, timing, and method of application are correct, but should NOT be applied directly to the crop if bees are present; H=highly toxic, severe losses expected, -- = data not available.
7. In accordance with the USDA National Organic Program, the Organic Materials Research Institute (OMRI) maintains a directory of all products that OMRI has determined are allowed for use in organic production, processing, and handling. These products are catalogued online in the **OMRI Products List** (see <https://www.omri.org/omri-lists>).

If you are having a **medical emergency** after using pesticides, always **call 911** immediately.



In Case of an Accident

- Remove the person from exposure
- Get away from the treated or contaminated area immediately
- Remove contaminated clothing
- Wash with soap and clean water
- Call a physician and/or the National Poison Control Center (1-800-222-1222).
Your call will be routed to your State Poison Control Center.
- **Have the pesticide label with you!**
- Be prepared to give the **EPA registration number** to the responding center/agency

Eggplant

Recommended Varieties¹

Type	Variety ^{1,2}	Days ³	F ₁ ⁴	Color	Calyx Color	Shape	Type	TMV ⁵
Standard Market Type	Aretussa RZ	65	Yes	White	Green	Elongated Oval		
	Gaudi	75	Yes	Black	Green	Oval Long		
	Nadia	70	Yes	Black	Green	Oval Long		X
	Night Shadow	68-75	Yes	Black	Green	Teardrop		
	Picasso	65	Yes	Purple/Black	Green	Teardrop		
	Santana	80	Yes	Black/Purple	Green	Elongated Oval		
Specialty Types	Annina	67	Yes	Purple Variegated	Green	Teardrop		
	Barbarella	65	Yes	Purple	Purple	Round	Sicilian	
	Calliope	64	Yes	Purple variegated	Green	Oval	Asian	
	Gretel	55	Yes	White	Green	Mini Slender	Japanese	
	Hansel	55	Yes	Purple	Green	Mini Slender	Japanese	
	Kermit	60	Yes	Green and White	Green	Mini Round	Thai	
	Megal	60	Yes	Purple/Black	Green	Elongated Oval	Italian	X
	Millionaire	55	Yes	Black	Purple	Slender	Japanese	
	Orient Express	58	Yes	Purple	Purple	Slender Long	Asian	
	Palermo	70	Yes	Purple	Purple	Round	Sicilian	
	Purple Shine	70	Yes	Purple	Purple	Slender Long	Chinese	
	Sabelle RZ	67	Yes	Purple	Purple	Round	Sicilian	
	Shoya Long	55-60	Yes	Purple	Purple	Slender Long	Japanese	

¹Listed alphabetically within type. ²Variety attributes based on Seed Company information. ³Days from transplanting till harvest. ⁴Hybrid (yes/no). ⁵TMV=Tobacco Mosaic Virus. Only those varieties with some resistance or tolerance to TMV are noted with an X.

Recommended Nutrients Based on Soil Tests

In addition to using the table below, check the suggestions on rate, timing, and placement of nutrients in your soil test report and Chapter B Soil and Nutrient Management. Your state's soil test report recommendations and/or your farm's nutrient management plan supersede the recommendations found below.

Eggplant ^{1,2}		Soil Phosphorus Level				Soil Potassium Level				Nutrient Timing and Method
		Low	Med	High (Opt)	Very High	Low	Med	High (Opt)	Very High	
	N (lb/A)	P ₂ O ₅ (lb/A)				K ₂ O (lb/A)				
	125-150 ³	250	150	100	0	250	150	100	0	Total nutrient recommended
	50-100	250	150	100	0	250	150	100	0	Broadcast and disk-in
	25-50	0	0	0	0	0	0	0	0	Sidedress 3-4 weeks after planting
	25-50	0	0	0	0	0	0	0	0	Sidedress 6-8 weeks after planting

For plasticulture, fertilization rates are based on a standard row spacing of 6 ft. ¹Apply 1-2 lb/A of boron (B) with broadcast fertilizer; see also Table B-7. in Chapter B Soil and Nutrient Management. ²Apply 25-30 lb/A of sulfur (S) for most soils. ³If crop is to be mulched with plastic but not drip/trickle fertilized, broadcast 125 lb/A N with recommended P₂O₅ and K₂O and disk-in or incorporate prior to laying mulch.

Plant Tissue Testing

Plant tissue testing can be a valuable tool to assess crop nutrient status during the growing season to aid with in-season fertility programs or to evaluate potential deficiencies or toxicities. Critical eggplant tissue test values for most recently matured leaves at early fruit set are N 4.2-6.0 %, P 0.3-0.7 %, K 3.5-5.0 %, Ca 0.8-1.5%, Mg 0.25-0.6% and S 0.4-0.6%. For additional nutrients and other growth stages consult with a tissue testing laboratory or check the following University of Florida website: <https://edis.ifas.ufl.edu/publication/ep081>.

Seed Treatment

Use hot water seed treatment - see section E 4.3. Disease Control in Seeds, Plant Growing Mix, and Plant Beds.

Transplant Production and Transplanting Dates

Sow seeds in the greenhouse 8-10 weeks before field planting. Three to four ounces of seed are necessary to produce plants for 1 acre. Optimum temperatures for germination and growth are 70-75°F. Seedlings should be transplanted

F. Eggplant

to 2-inch or larger pots any time after the first true leaves appear, or seed can be sown directly into the pots and thinned to a single plant per pot.

Harden plants for a few days at 60-65°F and set in field after danger of frost when average daily temperatures have reached 65-70°F. Usual transplanting period is May 15 to June 5. Eggplant is a warm-season crop that grows best at temperatures between 70-85°F. Temperatures below 65°F result in poor growth and fruit set.

Spacing Rows: 4-5 feet apart; plants: 2-3 feet apart in the row. Space plants 18-30 inches apart in PA.

Drip/Trickle Fertilization

Before mulching, adjust soil pH to around 6.5 and then apply enough farm-grade fertilizer to supply 60 lb/A of N, P₂O₅ and K₂O. Thoroughly incorporate fertilizer into the soil. If soil tests medium or less in soil K, apply a fertilizer with a ratio of 1-1-2 or 1-1-3 carrying 60 lb/A of N. After mulching and installing the drip irrigation system, apply completely soluble fertilizers to supply 40 lb/A (10-20 lb/A in PA) of N, P₂O₅ and K₂O during each application. On soils testing low and low to medium in Boron (B) and that have not received any preplant B fertilizer, include 0.25 lb/A of actual B in each soluble fertilizer application. The first soluble fertilizer application should be applied through the trickle irrigation system within 1 week after field transplanting. The same rate of soluble fertilizer should be applied about every 3 weeks during the growing season for a total of 6-7 applications.

Mulching and Fumigation

The use of black plastic mulch can increase eggplant yield and promote earliness. Various widths of plastic are available for different production systems and available equipment. At least 50% of the N should be in nitrate form (NO₃⁻¹) when planting in fumigated soil under plastic mulch. For more details, see the Weed Control section below.

Staking

High intensity eggplant production can benefit from staking with increased yields and less branch breakage. Use a staking system similar to that described for tomatoes. Pruning is not required for eggplant but removing the two lowest branches helps with plastic removal at seasons end if the plants are mowed off.

Harvest and Post-Harvest Considerations

Fruit should be harvested when the skin is still a glossy color, and the seed and pulp are white. Soft fruit and dark seed indicate over maturity. Mature fruit must be harvested to ensure continued fruit set. Harvested fruit should be moved to a protected area as soon as possible. If left in direct sunlight the fruit will sunburn. Cool eggplants in a cold room with forced-air, or with forced-air and evaporative cooling. Fruit are sensitive to temperatures below 50°F (see fruit disorders below) but can be stored for 1-2 weeks at 50-54°F and 90-95% relative humidity.

Fruit Disorders

Liver Spot and Pitting: ‘Liver spot’ and ‘pitting’ are late season physiological disorders that become apparent on the fruit surface post-harvest. Light tan to copper-colored spots and scratching may appear after washing; scratching is most likely caused by rough handling or contact of fruit with the ground. Pitting (small slightly sunken brown pits) may also occur. Liver spot and/or pitting are thought to be caused by a thinner waxy fruit cuticle as a result of cooler temperatures. Temperatures at or below 50°F are often associated with both disorders.

Internal Seed Cavity Browning: Symptoms of internal seed cavity browning include the discoloration or browning of the fruit tissue directly surrounding the seed cavity. The discoloration can be caused by low temperatures and/or bruising and compression injury during harvest and post-harvest handling.

Weed Control

THE LABEL IS THE LAW-see the Pesticide Use Disclaimer on the first page of Chapter F.

Recommended Herbicides

1. Identify the weeds in each field and select recommended herbicides. More information is available in the “Herbicide Effectiveness on Common Weeds in Vegetables” (Table E-3) in Chapter E Pest Management.
2. Minimize herbicide resistance development. Identify the herbicide mode of action group number and follow recommended good management practices; **bolded group numbers in tables below are herbicides at higher risk for selecting resistant weed populations.** Include non-chemical weed control whenever possible.

Labeled Application Sites for Eggplant									
Herbicide (*=Restricted Use)	HRAC group number	Plastic mulch production					Bareground production		
		Soil-Applied		Postemergence			Soil- applied	POST	Post- harvest
		Under Plastic	Row Middles	Over Plastic	Row Middles	Post- Harvest			
Sandea	2		YES		YES		directed ¹		
Prowl H2O	3		YES				YES ²		
Prefar	8	YES	YES				YES		
Devrinol	15	YES	YES				YES		
Poast	1			YES				YES	
Select / Select Max Shadow 3EC	1			YES				YES	
Gramoxone* ⁴	22				YES	YES	YES ³		YES

¹Sandea is labeled for bareground only if the spray is directed to the row middles. ²Transplants only.

³Gramoxone can be applied early preplant, or after planting but before crop emergence.

⁴Supplemental Label, be sure it is registered for the specific state and for the intended use.

1. Pre-Transplant Over Plastic

Group	Product Name (*=Restricted Use)	Product Rate	Active Ingredient	Active Ingredient Rate	PHI (d)	REI (h)
22	Gramoxone SL 3.0*	1.3 to 2.7 pt/A	paraquat	0.5 to 1.0 lb/A	--	24

-Gramoxone can be used for preplant weed control over the top of plastic mulch. Sufficient rainfall or sprinkler irrigation is needed to wash off the Gramoxone prior to planting to prevent damage to the crop.
-Restricted-use pesticide. Only certified applicators, who successfully complete the paraquat-specific training, can mix, load, or apply paraquat. Application of paraquat "under the direct supervision" of a certified applicator is no longer allowed. Required training link (<https://campus.extension.org/enroll/index.php?id=2201>); certified applicators must repeat training every three years.
 -Do not exceed 8 pt/A per season.
 -Rainfastness is 30 min.

2. Soil-Applied

Group	Product Name (*=Restricted Use)	Product Rate	Active Ingredient	Active Ingredient Rate	PHI (d)	REI (h)
2	Sandea 75DF	0.5 to 1 oz/A	halosulfuron	0.023 to 0.047 lb/A	30	12

-Plasticulture: row middles only; adjust equipment to keep the spray off the plastic.
-Bareground: apply between rows of direct-seeded or transplants. **Do not** apply as broadcast application; avoid contact of the herbicide with the planted crop
 -Suppresses or controls yellow nutsedge and certain broadleaf weeds. Sandea provides both residual and postemergence control of susceptible weed species. Effective postemergence control requires an adjuvant.
 -Sandea is an ALS inhibiting herbicide and resistant weed populations are common in the region.
-Do not use Group 2 herbicides repeatedly in the same field.
Do not apply Sandea to crops treated with a soil applied organophosphate insecticide or use a foliar applied organophosphate insecticide within 21 days before or 7 days after a Sandea application.
 -Maximum Sandea applications per year is 2 and **do not** exceed 2 oz/A during the crop season.

3	Prowl H2O 3.8CS	1 to 3 pt/A	pendimethalin	0.48 to 1.42 lb/A	70	24
---	-----------------	-------------	----------------------	-------------------	----	----

-Plasticulture: recommended for row middles only. Labeled for under plastic, but no local data or experience with this application.
-Bareground: broadcast preplant or preplant incorporated before transplanting; not labeled for direct-seeded crop.
 -Avoid root contact with Prowl-treated soil when placing transplants into furrow or hole or injury may occur.
 -Prowl labeled for directed application to transplanted or established direct-seeded eggplant; avoid contact with leaves or stems.
 -Prowl will not control emerged weeds, only provides residual control; row middle applications may be made with Gramoxone using shielded sprayers. Use the lower rate on coarse-textured or sandy soils. Activate with ½ inch of rainfall or sprinkler irrigation within 48 hours of application to control most annual grasses and certain broadleaf weeds.
 -Maximum Prowl H2O application per season: 3 pt/A.

8	Prefar 4E	5 to 6 qt/A	bensulide	5 to 6 lb/A	--	12
---	-----------	-------------	------------------	-------------	----	----

-Plasticulture under plastic: apply in a band under the plastic, immediately before laying the mulch. Allow 7 days before making transplant holes to allow condensation to incorporate the herbicide. Plasticulture: row middles application is labeled.
-Bareground: apply preemergence or preplant incorporated.
-Do not incorporate more than 2 inches deep (1 inch is optimum). If applied preemergence, irrigate within 36 h of application with ½ inch of water; if not incorporated with irrigation or rainfall within 36 h, weed control maybe reduced.
 -Provides control/suppression of some annual grass weeds and some broadleaves including pigweeds, purslane, and lambsquarters.

2. Soil Applied - continued next page

F. Eggplant

2. Soil Applied - continued

15	Devrinol 2-XT 2EC Devrinol DF-XT 50DF	2 to 4 qt/A 2 to 4 lb/A	napropamide	1.0-2 lb/A	--	24
----	--	----------------------------	--------------------	------------	----	----

-**Plasticulture**: labeled for under plastic mulch; apply in a band under the plastic, immediately before laying mulch. Condensation that forms on the underside of the mulch will activate the herbicide. Plasticulture: row middles application is labeled.
 -**Bareground**: apply as broadcast, preemergence treatment for transplanted eggplant. Rainfall or irrigation within 24 h after application improves performance (½ inch sprinkler irrigation).
 -Annual grasses and certain annual broadleaf weeds will be suppressed or controlled. May reduce stand and yield of fall planted small grain crop. Moldboard plowing will reduce the risk of injury. -Maximum application per season: 4 qt/A (2-XT) or 4 lb/A (DF-XT).

3. Postemergence

Group	Product Name (*= Restricted Use)	Product Rate	Active Ingredient	Active Ingredient Rate	PHI (d)	REI (h)
1	Shadow 3EC	4 to 5.33 fl oz/A	clethodim	0.07 to 0.125 lb/A	20	24
	Select 2EC	6 to 8 fl oz/A				
	Select Max 0.97EC	9 to 16 fl oz/A				
	Poast 1.5EC	1 to 1.5 pt/A	sethoxydim	0.2 to 0.3 lb/A	20	12

-**Select 2EC**: use crop oil concentrate (COC) at 1% v/v (1 gal/100 gal of spray solution). **Select Max 0.97EC**: use nonionic surfactant (NIS) at 0.25% v/v (1 qt/100 gal of spray solution). **Shadow 3EC**: use crop oil concentrate (COC) at 1% v/v (1 gal/100 gal of spray solution) for large or stressed grasses; use nonionic surfactant (NIS) at 0.25% v/v (1 qt/100 gal of spray solution) when crop safety is a concern. **Poast 1.5EC**: Use COC at 1.0% v/v.
 -**General comments**: -The use of COC may increase the risk of crop injury when hot or humid conditions prevail. To reduce the risk of crop injury, switch to NIS when grasses are small and soil moisture is adequate. -Use lower labeled rates for annual grass control and higher labeled rates for perennial grass control. For best results, treat annual grasses when they are actively growing and before tillers are present. -Yellow nutsedge, wild onion, wild garlic, and broadleaf weeds will **not** be controlled with these herbicides. -These herbicides control many annual and certain perennial grasses. Clethodim is best on annual bluegrass; while Poast is preferred for goosegrass control. -Repeated applications may be necessary to control certain perennial grasses. If repeat applications are necessary, allow 14 days between applications. -Rainfastness is 1 h.
 -**Do not** tank mix with or apply within 2 to 3 days of any other pesticide, unless labeled, as this may increase the risk of crop injury or reduce the control of grasses. -**Do not** apply more than 8 fl oz/A of Select 2EC in a single application and **do not** exceed 2 pt/A for the season; **do not** apply more than 16 fl oz/A of Select Max in a single application and **do not** exceed 4 pt/A for the season.
 -**Do not** apply more than 5.33 fl oz/A of Shadow 3EC in a single application and **do not** exceed 21.33 fl oz/A for the season.
 -**Do not** apply more than 1.5 pt/A Poast 1.5EC in a single application and **do not** exceed 4.5 pt/A for the season.

22	Gramoxone SL 3.0*	1.3 pt/A	paraquat	0.5 lb/A	--	24
----	-------------------	----------	-----------------	----------	----	----

-Gramoxone can be applied before or after seeding to control emerged broadleaf weeds and grass seedlings.
 -For use in plasticulture: row middles as a shielded application.
 -Include a nonionic surfactant at 0.25% v/v. **Do not** allow spray to contact crop foliage as injury may result. Use flaps that drag along the edge of plastic mulch and use low spray pressure (maximum of 30 psi) to reduce small droplets that are prone to drift.
 -See the label for additional information and warnings. Rainfastness is 30 min. A maximum of 3 applications per year are allowed.
 -**Restricted-use pesticide**. Only certified applicators, who successfully complete the paraquat-specific training, can mix, load, or apply paraquat. Application of paraquat "under the direct supervision" of a certified applicator is no longer allowed. Required training link (<https://campus.extension.org/enrol/index.php?id=2201>); certified applicators must repeat training every three years.

4. Postharvest

Group	Product Name (*= Restricted Use)	Product Rate	Active Ingredient	Active Ingredient Rate	PHI (d)	REI (h)
22	Gramoxone SL 3.0*	1.5 to 2 pt/A	paraquat	0.56 to 0.75 lb/A	--	24

-**Supplemental Label in DE for postharvest application to desiccate the crop**. -Apply after the last harvest for bareground or plasticulture. Always include an adjuvant. -Spray coverage is essential for optimum effectiveness. See the label for additional information and warnings. -Rainfastness 30 min. A maximum of 2 applications for crop desiccation are allowed. -**Restricted-use pesticide**. Only certified applicators, who successfully complete the paraquat-specific training, can mix, load, or apply paraquat. Application of paraquat "under the direct supervision" of a certified applicator is no longer allowed. Required training link (<https://campus.extension.org/enrol/index.php?id=2201>); certified applicators must repeat training every three years.

5. Other Labeled Herbicides

These products are labeled but limited local data are available; and/or are labeled but not recommended in our region due to potential crop injury concerns.

Group	Product Name (*= Restricted Use)	Active Ingredient
14	Aim (hooded or directed application only)	carfentrazone
14	Vida	pyraflufen
14	Aquestra, others	sulfentrazone

Insect Control

THE LABEL IS THE LAW-see the Pesticide Use Disclaimer on the first page of Chapter F. Recommended Insecticides

Note: For **premixes**, the group number (representing the mode of action) and active ingredient that contributes the most to control is generally listed first. In some cases, only one ingredient in a premix provides control.

Aphids

Green peach aphids (GPA) are the most common aphids on eggplant. Winged females can produce numerous live pale, yellow or pink-colored young (nymphs). Tremendous numbers can build up on the undersides of leaves often following pyrethroid insecticide applications. Aphids are sucking insects. They excrete a sugary, sticky substance (“honeydew”) that can cause growth of black sooty mold fungus. Both honeydew and mold on fruit can hurt its marketability. Predators and parasitoids (braconid wasps) often can keep aphid populations below damaging levels. Broad-spectrum insecticides, like pyrethroids, destroy these natural enemies. Use selective insecticides whenever possible. Sample plants for aphids as well as the presence of natural enemy species. Spray only when aphid densities appear to be increasing in the absence of predators.

Apply one of the following formulations (note: spray coverage to the underside of the leaf is important):						
Group	Product Name (* = Restricted Use)	Product Rate	Active Ingredient(s)	PHI (d)	REI (h)	Bee TR
1A	Lannate LV*	0.75 to 3.0 pt/A	methomyl - GPA only	5	48	H
1B	Malathion 57 EC	1 to 1.5 pt/A	malathion	3	12	H
4A	Neonicotinoid insecticides registered for use on eggplant: see table at the end of insect control.					
4C	Transform WG	0.75 to 1.0 oz/A	sulfoxaflor	1	24	H
4C + 3A	Ridgeback*	4.5 to 13.8 fl oz/A	sulfoxaflor + bifenthrin	7	24	H
4D	Sivanto Prime	21.0 to 28.0 fl oz/A	flupyradifurone - soil	45	4	M
4D	Sivanto Prime	7.0 to 14.0 fl oz/A	flupyradifurone - foliar	1	4	M
9B	Fulfill 50WDG	2.75 oz/A	pymetrozine	0	12	L
9B	PQZ	2.4 to 3.2 fl oz/A	pyrifluquinazon	1	12	L
9D	Sefina	3.0 fl oz/A	afidopyropen	0	12	L
21A	Torac	17.0 to 21.0 fl oz/A	tolfenpyrad	1	12	H
23	Boxadon 360	2.1 to 3.4 fl oz/A	spirotetramat	1	24	L
23	Movento	4.0 to 5.0 fl oz/A	spirotetramat	1	24	L
23 + 7C	Senstar	8 to 10 fl oz/A	spirotetramat + pyriproxyfen	1	24	L
28 + 6	Minecto Pro* ¹	10.0 fl oz/A	cyantraniliprole + abamectin	7	12	H
29	Beleaf 50SG	2.8 to 4.28 oz/A	flonicamid	0	12	L
UNF	BotaniGard ES	0.25 to 1 qt/A	<i>Beauveria bassiana</i> , strain GHA	0	4	L

¹Use of a non-sticker adjuvant is required.

Colorado Potato Beetles (CPB)

CPB rapidly develops resistance to insecticides (see also section E 3.2. Insecticide Mode of Action: Reducing the Risk of Insecticide Resistance Development). Augmentative releases of the egg parasitoid, *Edovum puttleri*, has been shown to control CPB effectively in eggplant, or apply one of the following insecticides.

Apply one of the following formulations:						
Group	Product Name (* = Restricted Use)	Product Rate	Active Ingredient(s)	PHI (d)	REI (h)	Bee TR
1A	Vydate L*	2.0 to 4.0 pt/A	oxamyl – foliar	1	48	H
4A	Neonicotinoid insecticides registered for use on Eggplant: see table at the end of Insect Control.					
4D	Sivanto Prime	10.5 to 14.0 fl oz/A	flupyradifurone - foliar	1	4	M
5	Entrust SC (OMRI)	3.0 to 6.0 fl oz/A	spinosad	1	4	H
5	Radiant SC	5.0 to 10.0 fl oz/A	spinetoram	1	4	H
6	Agri-Mek SC* ¹	1.75 to 3.5 fl oz/A	abamectin	7	12	H
11A	Trident (OMRI) ²	3.0 to 6.0 qt/A	<i>Bacillus thuringiensis tenebrionis</i> ¹	0	4	L
15	Rimon 0.83EC	9.0 to 12.0 fl oz/A	novaluron	1	12	M
15 + 4A	Cormoran	9.0 fl oz/A	novaluron + acetamiprid	7	12	M
21A	Torac	14.0 to 21.0 fl oz/A	tolfenpyrad	1	12	H
28	Coragen 1.67SC	3.5 to 7.5 fl oz/A	chlorantraniliprole	1	4	L
	Coragen eVo	1.2 to 2.5 fl oz/A				

Colorado Potato Beetles (CPB) - continued next page

F. Eggplant

Colorado Potato Beetles (CPB) - continued

Group	Product Name	Product Rate	Active Ingredient(s)	PHI (d)	REI (h)	Bee TR
28	Exirel	7.0 to 13.5 fl oz/A	cyantraniliprole	1	12	H
28	Harvanta 50SL	10.9 to 16.4 fl oz/A	cyfluprol	1	4	H
28	Verimark	5.0 to 10.0 fl oz/A	cyantraniliprole - soil (drip or injection)	1	4	H
28 + 3A	Besiege*	6.0 to 9.0 fl oz/A	chlorantraniliprole + lambda cyhalothrin	5	24	H
28 + 3A	Elevest*	7.0 to 9.6 fl oz/A	chlorantraniliprole + bifenthrin	7	12	H
28 + 6	Minecto Pro* ¹	5.5 to 10.0 fl oz/A	cyantraniliprole + abamectin	7	12	H
30	Incipio	2.1 to 4.1 fl oz/A	isocycloseram	1	12	H

¹Use of a non-sticker adjuvant is required. ²Larval reduction may not be noticeable for 48-72 h. Apply when eggs begin to hatch and repeat at 5-7-day intervals. If rainfall occurs within 24 h post-treatment, reapplication may be necessary.

Eggplant Lacebugs

Eggplant lacebug is a small sucking insect with lacy wings and conspicuous veins. It can cause stippling and yellowing/whitening of leaves. Most insecticides are not labeled for this sporadic pest; however, use of any insecticide labeled for flea beetles will provide adequate control of this pest. Good insecticide coverage is essential.

Flea Beetles

Apply one of the following formulations:						
Group	Product Name (* = Restricted Use)	Product Rate	Active Ingredient(s)	PHI (d)	REI (h)	Bee TR
3A	Pyrethroid insecticides registered for use on Eggplant: see table at the end of Insect Control.					
4A	Neonicotinoid insecticides registered for use on Eggplant: see table at the end of Insect Control.					
5	Entrust SC (OMRI)	4.0 to 8.0 fl oz/A	spinosad	1	4	H
21A	Torac	17.0 to 21.0 fl oz/A	tolfenpyrad	1	12	H
28	Verimark	6.75 to 13.5 fl oz/A	cyantraniliprole - soil at planting	1	4	H
30	Incipio	2.1 to 4.1 fl oz/A	isocycloseram	1	12	H
UNK	BotaniGard ES (OMRI)	0.25 to 1 qt/A	<i>Beauveria bassiana</i> , strain GHA	0	4	L

Leafminers

Apply one of the following formulations:						
Group	Product Name (* = Restricted Use)	Product Rate	Active Ingredient(s)	PHI (d)	REI (h)	Bee TR
4A	Neonicotinoid insecticides registered for use on Eggplant: see table at the end of Insect Control.					
5	Entrust SC (OMRI)	6.0 to 10.0 fl oz/A	spinosad	1	4	H
5	Radiant SC	6.0 to 10.0 fl oz/A	spinetoram	1	4	H
6	Agri-Mek SC* ¹	1.75 to 3.5 fl oz/A	abamectin	7	12	H
15	Rimon 0.83EC	12 fl oz/A	novaluron	1	12	M
15 + 4A	Cormoran	12.0 fl oz/A	novaluron + acetamiprid	7	12	M
28	Exirel	13.5 to 20.5 fl oz/A	cyantraniliprole	1	12	H
28	Harvanta 50SL	10.9 to 16.4 fl oz/A	cyfluprol	1	4	H
28	Verimark	6.75 to 13.5 fl oz/A 6.75 to 10.0 fl oz/A	cyantraniliprole - soil at planting cyantraniliprole - soil (drip or injection)	1	4	H
28 + 6	Minecto Pro* ¹	5.5 to 10.0 fl oz/A	cyantraniliprole + abamectin	7	12	H
30	Incipio	2.1 to 4.1 fl oz/A	isocycloseram	1	12	H

¹Use of a non-sticker adjuvant is required.

Mites

Apply one of the following formulations:						
Group	Product Name (* = Restricted Use)	Product Rate	Active Ingredient(s)	PHI (d)	REI (h)	Bee TR
6	Agri-Mek SC* ¹	1.75 to 3.5 fl oz/A	abamectin	7	12	H
6 + 28	Minecto Pro* ¹	5.5 to 10.0 fl oz/A	abamectin + cyantraniliprole	7	12	H
10A	Onager 1EC	12 to 24 fl oz/A	hexythiazox	1	12	N
10B	Zeal Miticide	2.0 to 3.0 oz/A	etoxazole	7	12	L
10B	Zeal MVP	23.0 to 34.6 fl oz/A	etoxazole	7	12	L
12B	Vendex 50WP*	2.0 to 3.0 lb/A	fenbutatin-oxide	3	48	N
20B	Kanemite 15SC	31 fl oz/A	acequinocyl	1	12	L
20D	Acramite 50WS	0.75 to 1.0 lb/A	bifenazate	3	12	M

Mites - continued next page

Mites - continued

21A	Magister SC	24.0 to 36.0 fl oz/A	fenazaquin	3	12	H
21A	Portal	2.0 pt/A	fenpyroximate	1	12	L
21A	Torac	14.0 to 21.0 fl oz/A	tolfenpyrad - broad mite only	1	12	H
23	Boxadon 360	2.1 to 3.4 fl oz/A	spirotetramat - broad mite and tomato russet mite	1	24	L
23	Movento	4.0 to 5.0 fl oz/A	spirotetramat - broad mite and tomato russet mite	1	24	L
23	Oberon 2SC	8.5 fl oz/A	spiromesifen	1	12	M
30	Incipio	2.1 to 4.1 fl oz/A	isocycloseram	1	12	H

¹Use of a non-sticker adjuvant is required.

Thrips

Apply one of the following formulations:						
Group	Product Name (* = Restricted Use)	Product Rate	Active Ingredient(s)	PHI (d)	REI (h)	Bee TR
3A ¹	Pyrethroid insecticides registered for use on Eggplant: see table at the end of Insect Control.					
4A ²	Neonicotinoid insecticides registered for use on Eggplant: see table at the end of Insect Control.					
5	Entrust SC (OMRI)	4.0 to 8.0 fl oz/A	spinosad	1	4	H
5	Radiant SC	6.0 to 10.0 fl oz/A	spinetoram	1	4	H
21A	Torac	21.0 fl oz/A	tolfenpyrad	1	12	H
30	Incipio	3.1 to 5.1 fl oz/A	isocycloseram	1	12	H

¹Resistance concerns with western flower thrips. ²Resistance concerns with tobacco thrips.

[Insecticides with Suppression Only on the label: Movento, Boxadon 360, Harvanta, Verimark]

Group 3A Pyrethroid Insecticides Registered for Use on Eggplant						
Apply one of the following formulations (check if the product label lists the insect you intend to spray; the label is the law). Note: resistance concerns with western flower thrips.						
Product Name (* = Restricted Use)	Product Rate	Active Ingredient(s)	PHI (d)	REI (h)	Bee TR	
Asana XL*	5.8 to 9.6 fl oz/A	esfenvalerate	7	12	H	
Baythroid XL*	1.6 to 2.8 fl oz/A	beta-cyfluthrin	7	12	H	
Brigade 2EC*, Brigade eVo	2.1 to 6.4 fl oz/A	bifenthrin	7	12	H	
Declare*	0.77 to 1.54 fl oz/A	gamma-cyhalothrin	5	24	H	
Fastac CS	2.2 to 3.8 fl oz/A	alpha-cypermethrin	1	12	H	
Hero*	4.0 to 10.3 fl oz/A	zeta-cypermethrin + bifenthrin	7	12	H	
Lambda-Cy EC*, others	1.92 to 3.84 fl oz/A	lambda-cyhalothrin	5	24	H	
Mustang Maxx*	2.24 to 4.0 fl oz/A	zeta-cypermethrin	1	12	H	
Permethrin 3.2EC*, others	4.0 to 6.0 fl oz/A	permethrin	3	12	H	
PyGanic Crop protection EC 5.0 II (OMRI)	4.5 to 15.61 fl oz/A	pyrethrins	0	12	H	
Tombstone*	1.6 to 2.8 fl oz/A	cyfluthrin	7	12	H	
Warrior II*	1.28 to 1.92 fl oz/A	lambda-cyhalothrin	5	24	H	
Combo products containing a pyrethroid						
Besiege*	6.0 to 9.0 fl oz/A	lambda-cyhalothrin + chlorantraniliprole (Group 28)	5	24	H	
Brigadier*	3.8 to 9.85 fl oz/A	bifenthrin + imidacloprid (Group 4A) - foliar	7	12	H	
Elevest*	5.6 to 9.6 fl oz/A	bifenthrin + chlorantraniliprole (Group 28)	7	12	H	
Endigo ZC* and ZCX*	4.0 to 4.5 fl oz/A	lambda-cyhalothrin + thiamethoxam (Group 4A)	5	24	H	
Leverage 360*	3.8 to 4.1 fl oz/A	beta-cyfluthrin + imidacloprid (Group 4A)	7	12	H	
Ridgeback*	4.5 to 13.8 fl oz/A	bifenthrin + sulfoxaflor (Group 4C)	7	24	H	

Group 4A Neonicotinoid Insecticides Registered for Use on Eggplant						
Apply one of the following formulations (check if the product label lists the insect you intend to spray; the label is the law). Note: resistance concerns with tobacco thrips.						
Product Name (* = Restricted Use)	Product Rate	Active Ingredient(s)	PHI (d)	REI (h)	Bee TR	
Actara 25WDG	2.0 to 5.5 oz/A	thiamethoxam	0	12	H	

Group 4A Neonicotinoid Insecticides Registered for Use on Eggplant - continued next page

F. Eggplant

Group 4A Neonicotinoid Insecticides Registered for Use on Eggplant - continued

Admire Pro	7.0 to 10.5 fl oz/A	imidacloprid - soil	21	12	H
Admire Pro	1.3 to 2.2 fl oz/A	imidacloprid - foliar	21	12	H
Assail 30SG	1.5 to 4.0 oz/A	acetamiprid	7	12	M
Assail 30SC	1.3 to 3.4 fl oz/A	acetamiprid	7	12	M
Platinum 75SG	1.66 to 3.67 oz/A	thiamethoxam	30	12	H
Scorpion 35SL	9.0 to 13 fl oz/A	dinotefuran - soil	21	12	H
Scorpion 35SL	2.0 to 7.0 fl oz/A	dinotefuran - foliar	21	12	H
Venom	5.0 to 7.5 oz/A	dinotefuran - soil	21	12	H
Venom	1.0 to 4.0 oz/A	dinotefuran - foliar	21	12	H
Combo products containing a neonicotinoid					
Brigadier*	3.80 to 9.85 fl oz/A	imidacloprid + bifenthrin (Group 3A) - foliar	7	12	H
Cormoran	9.0 to 12.0 fl oz/A	acetamiprid + novaluron (Group 15)	7	12	M
Durivo	10.0 to 13.0 fl oz/A	thiamethoxam + chlorantraniliprole (Group 28)	30	12	H
Endigo ZC* and ZCX*	4.0 to 4.5 fl oz/A	thiamethoxam + lambda-cyhalothrin (Group 3A)	5	24	H
Leverage 360*	3.8 to 4.1 fl oz/A	imidacloprid + beta-cyfluthrin (Group 3A)	7	12	H
Voliam Flexi	4.0 to 7.0 oz/A	thiamethoxam + chlorantraniliprole (Group 28)	1	12	H

Disease Control

THE LABEL IS THE LAW-see the Pesticide Use Disclaimer on the first page of Chapter F. Recommended Fungicides

Nematodes

Salibro is labeled at 30.7 to 61.4 fl oz/A pre-plant incorporated, pre-plant drip, or at-plant drip. In-season drip application rates are 15.4 to 30.7 fl oz/A. See sections E 1.5. Soil Fumigation and E 1.6. Nematode Control.

Seed Treatment

Use hot water seed treatment - see section E 4.3. Disease Control in Seeds, Plant Growing Mix, and Plant Beds.

Damping-off caused by *Phytophthora*, *Pythium*, and *Rhizoctonia*

Code	Product Name (*=Restricted Use)	Product Rate	Active Ingredient(s)	PHI (d)	REI (h)	Bee TR
Apply one of the following at-planting (see label for application methods and restrictions):						
Phytophthora and Pythium root rot¹						
4	Ridomil Gold 4SL	0.5 to 1.0 pt/A	mefenoxam	5	48	N
4	Ultra Flourish 2E	2.0 to 4.0 pt/A	mefenoxam	5	48	N
4	MetaStar 2E AG	4.0 to 8.0 pt/A	metalaxyl	AP	48	N
4 + 49	Orondis Gold ²	28.0 to 55.0 fl oz/A	mefenoxam + oxathiapiprolin	AP	48	--
21	Ranman 400SC ²	2.75 fl oz/A	cyazofamid	AP	12	L
Phytophthora, Pythium, and Rhizoctonia root rot						
4 + 11	Uniform 3.72SC	0.34 fl oz/1000 ft row. Avoid direct seed contact, which may cause delayed emergence.	mefenoxam + azoxystrobin	AP	0	N
Rhizoctonia root and stem rot						
11	azoxystrobin 2.08F ³	0.40 to 0.80 fl oz/1000 ft row	azoxystrobin	AP	4	N
3 + 7	Aprovia Top 1.62EC ⁴	10.5 to 13.5 fl oz/A	difenoconazole + benzovindiflupyr	14	12	--

¹Also see Phytophthora Blight - Root and Crown Rot below.

²Orondis Gold and Ranman can be applied in transplant water. See label for specifics.

³Rhizoctonia can become a problem in transplants that have been in transplant trays for too long prior to transplanting, or in transplants shortly after planting where the root zone is allowed to become excessively dry. To help suppress Rhizoctonia root rot apply the following via drip at transplanting.

⁴Apply as a foliar application for bare soil beds; will also help suppress Southern Blight

Bacterial and Fungal Diseases

Phytophthora Blight (*Phytophthora capsici*) - Root and Crown Rot

To minimize the occurrence of Phytophthora blight, rotate fields away from susceptible crops (such as cucurbits, peppers, eggplants, and tomatoes) for as many years as possible. Avoid using mefenoxam if insensitivity is known to exist. Sensitivity to mefenoxam can return if it has not been used in recent years.

Code	Product Name (* = Restricted Use)	Product Rate	Active Ingredient(s)	PHI (d)	REI (h)	Bee TR
Apply one of the following formulations via drip application at transplanting and 30 days later:						
4	Ridomil Gold 4SL	1.0 pt/A	mefenoxam	7	12	N
4	Ultra Flourish 2E	1.0 qt/A	mefenoxam	7	12	N
4 + 49	Orondis Gold ^{1,2}	28.0 to 55.0 fl oz/A	mefenoxam + oxathiapiprolin	7	48	--
If conditions favor disease development, apply the following drip application 14 d after at-transplanting applications:						
43	Presidio 4SC	3.0 to 4.0 fl oz/A	fluopicolide	2	12	L

¹If Orondis Gold is applied via drip application it cannot be applied as a foliar spray.

²Orondis Gold can be applied in transplant water, see label for rates and restrictions.

Phytophthora Blight (*Phytophthora capsici*) - Fruit and Stem Rot

Code	Product Name (* = Restricted Use)	Product Rate	Active Ingredient(s)	PHI (d)	REI (h)	Bee TR
For suppression of the aerial stem and fruit rot phase of Phytophthora blight, apply and rotate the following with a fixed copper at labeled rates on a 7 to 10 day schedule or when environmental conditions are conducive for disease development:						
21	Ranman 400SC	2.75 fl oz/A PLUS a non-ionic surfactant (do not apply Ranman with copper)	cyazofamid	0	12	L
40	Forum 4.17SC	6.0 fl oz/A	dimethomorph	0	12	N
43	Presidio 4SC	3.0 to 4.0 fl oz/A	fluopicolide	1	12	L
40+49	Orondis Ultra 2.33SC	5.5 to 8.0 fl oz/A	mandipropamid + oxathiapiprolin	1	4	--
4 + 49	Orondis Gold	28.0 to 55.0 fl oz/A	mefenoxam + oxathiapiprolin	0	4	--

¹If Orondis Gold is applied via a foliar application it cannot be applied via drip system. See label for restrictions.

Fungal Fruit Rots

Scout regularly and begin preventative sprays when weather conditions favor disease development and repeat every 7-10 days. Do not apply FRAC code 11 fungicides more than 4 times in a single year. Tank mix and rotate with a protectant fungicide such as fixed copper or chlorothalonil and rotate with other FRAC codes to help reduce resistance development.

Code	Product Name (* = Restricted Use)	Product Rate	Active Ingredient(s)	PHI (d)	REI (h)	Bee TR
Tank mix chlorothalonil at 1.5 pt 6F/A or fixed copper at labeled rates with one of the following FRAC code 11 fungicides:						
3 + 7	Aprovia Top 1.62EC	10.5 to 13.5 fl oz/A	difenoconazole + benzovindiflupyr	14	1	--
3 + 11	Topguard EQ 4.29SC	4.0 to 8.0 fl oz/A	flutriafol + azoxystrobin	0	1	--
3 + 11	Quadris Top 1.67SC	8.0 to 14.0 fl oz/A	difenoconazole + azoxystrobin	0	12	--
7 + 11	Priaxor 4.17SC	4.0 to 8.0 fl oz/A	fluxapyroxad + pyraclostrobin	7	12	N
And rotate with one of the following:						
M01	copper (OMRI) ¹	at labeled rates	copper	0	24	N
M05	chlorothalonil 6F	1.5 pt/A	chlorothalonil	3	12	M

¹There are several OMRI listed copper-based products; see labels for specifics. Copper applications may help suppress some fungal pathogens in organic production systems.

Fungal Leaf Spots

Scout on a regular basis and begin preventative sprays when weather conditions favor disease development, or when symptoms of disease first appear, and repeat every 7-10 days. Do not apply FRAC code 11 fungicides more than 4 times in a single year. Tank mix FRAC code 7 or 11 fungicides with a protectant fungicide and rotate with other FRAC codes to help reduce resistance development.

Code	Product Name (* = Restricted Use)	Product Rate	Active Ingredient(s)	PHI (d)	REI (h)	Bee TR
Tank mix chlorothalonil 6F at 1.5 pt/A or fixed copper at labeled rates and rotate among the following fungicides with different FRAC codes:						
7	Endura 70WG	2.5 to 3.5 oz/A	boscalid	0	12	--

Fungal Leaf Spots - continued next page

F. Eggplant

Fungal Leaf Spots - continued

7	Fontelis 1.67SC	10.0 to 24.0 fl oz/A	penthiopyrad	0	12	L
7 + 3	Luna Flex 3.13SC	8.0 to 13.6 fl oz/A	fludioxonil + difenoconazole	0	12	--
7 + 11	Luna Sensation 4.25SC	5.0 to 7.6 fl oz/A	fluopyram + trifloxystrobin	7	12	--
7 + 12	Miravis Prime	9.2 to 11.4 fl oz/A	pydiflumetofen + fludioxonil	0	12	--
11	azoxystrobin 2.08F	6.0 to 15.5 fl oz/A	azoxystrobin	0	4	N
11	Cabrio 20EG	8.0 to 12.0 oz/A	pyraclostrobin	0	12	N
3 + 11	Quadris Top 1.67SC	8.0 to 14.0 fl oz/A	difenoconazole + azoxystrobin	0	12	--
And rotate with one of the following:						
M01	copper (OMRI) ¹	at labeled rates	copper	0	24	N
M05	chlorothalonil 6F	1.5 pt/A	chlorothalonil	3	12	M
M05+22	Zing! 4.9SC	34 fl oz/A	chlorothalonil + zoxamide	3	12	M

¹There are several OMRI listed copper-based products; see labels for specifics. Copper applications may help suppress some fungal pathogens in organic production systems.

Verticillium Wilt

The best control can be accomplished by using a 4- to-5-year rotation with crops other than tomato, potato, pepper, strawberry, or any of the brambles. Varieties which appear to maintain yield in infested fields include Classic, and Epic. Soil fumigation will provide some control by delaying symptom expression. Use metam-sodium (Vapam HL - see label for specifics and restrictions). Broadcast treatments are superior to row treatments. Refer to section E 1.5. Soil Fumigation for details on application.

Viruses

Tomato Spotted Wilt Virus

Tomato Spotted Wilt Virus is spread by thrips from flowering ornamental plants to eggplant. Do not grow any ornamental bedding plants in the same greenhouse as eggplant transplants. Monitor and scout greenhouses for thrips and begin an insecticide control program once observed.