

F. Commodity Recommendations

Pesticide Use Disclaimer

THE LABEL IS THE LAW

Before using a pesticide, check the label for up to date rates and restrictions.

Labels can be downloaded from: <http://www.cdms.net/>, <https://www.greenbook.net/> or <http://www.agrian.com/labelcenter/results.cfm>

For more information on Pesticide Safety and the Pesticide Label see chapter D.

Guide to the Recommended Pesticide Tables in the Following Crop Sections:

- 1. Pesticides are listed by group or code number based on chemical structure and mechanism of action**, as classified by the Weed Science Society of America (WSSA) for herbicides, the Insecticide Resistance Action Committee (IRAC) for insecticides, and the Fungicide Resistance Action Committee (FRAC) for fungicides.
If the number is in bold font, the product may have resistance concerns.
- 2. For restricted use pesticides**, the restricted active ingredients are labeled with a *. (See section D 3.2.1 “Restricted Use Classification Statement” for more information).
- 3. In addition to the pesticides listed below, other formulations or brands with the same active ingredient(s) may be available. ALWAYS CHECK THE LABEL:**
 - a) to ensure a pesticide is labeled for the same use,**
 - b) to ensure the pesticide is labeled for the desired crop, and**
 - c) for additional restrictions.**
- 4. All pesticide recommendations are made for spraying a broadcast area of 1 acre** (43,560 square feet). **Adjust the rate for banded applications** (for more information, see section E 1.3 Calibrating Granular Applicators).
- 5. Check the label for the maximum amount of pesticide per application and the maximum number of applications per year.**
- 6. Bee Toxicity Rating (Bee TR):** N=nontoxic; L=minimum impact on bees; M=moderately toxic, can be used if dosage, timing and method of application are correct, but should NOT be applied directly to the crop if bees are present; H=highly toxic, severe losses expected, -- = data not available.

Cucumbers

For earlier cucumber production and higher, more concentrated yields, use gynoecious varieties. A gynoecious plant produces a high percentage of female flowers and fruit. To produce pollen, 1 to 15% of pollinator must be planted and seed companies add this seed to the gynoecious variety. Both pickling and slicing gynoecious varieties are available. Parthenocarpic cucumbers that produce fruit without pollination are also available for protected culture and field production.

Recommended Varieties¹

Type	Variety	Days	F1 ²	Type ³	Use ⁴	Reported Disease Resistance ⁵								
						Scab (Ccu)	PM (Px)	AN (Co)	DM ⁵ (Pcu)	ALS (Psl)	Cmv	Wmv	Zmv	Prsv
Standard Slicing Varieties	Bristol	54	Yes	Gyn	F	X	X	X		X	X	X	X	X
	Dasher II	58	Yes	Gyn	F	X	X	X		X	X			
	Dominator	55	Yes	Gyn	F	X	X	X		X	X			
	General Lee	66	Yes	Gyn	F	X	X			X				
	Intimidator	53	Yes	Gyn	F	X	X	X		X	X			
	Mongoose	55	Yes	Gyn	F	X	X	X		X	X	X	X	X
	Python	55	Yes	Gyn	F	X	X	X		X	X			
	Speedway	56	Yes	Gyn	F	X	X	X		X	X			
	Stonewall	53	Yes	Gyn	F	X	X	X		X	X			
	SV4719CS	56	Yes	Gyn	F	X	X	X	X	X			X	
	Talladega	61	Yes	Gyn	F	X	X	X		X	X	X		
Thunder	58	Yes	Gyn	F	X	X	X		X	X		X		
Slicers Long Types	Suyo Long	61	No	Mon	F		X							
	Tasty Green	52	Yes	Mon	F		X							
Pickles	Bowie	51	Yes	Parth	MP	X	X							
	Citadel	52	Yes	Gyn	HMP	X	X	X	X	X	X			
	Eureka	57	Yes	Mon	HF	X	X	X		X	X	X		X
	Expedition	50	Yes	Gyn	MP	X	X	X		X	X			
	Feisty	57	Yes	Gyn	MP	X	X	X		X	X			
	Jackson Supr.	52	Yes	Gyn	HMFP	X	X	X		X	X			
	Lafayette	52	Yes	Gyn	MP	X	X	X		X	X			
	Liszt	51	Yes	Parth	MP	X	X							
	Logan	51	Yes	Gyn	MP	X	X	X		X	X			
	Max Pack	57	Yes	Mon	FH	X	X	X		X	X	X	X	X
	NQ5007	50	Yes	Parth	MP	X	X	X		X	X			
	NQ5543	49	Yes	Parth	MP	X	X	X		X	X			
	Peacemaker	52	Yes	Gyn	MHP	X	X	X	X	X	X			
	Puccini	50	Yes	Parth	HMFP	X	X	X		X	X			
	Rubenstein	51	Yes	Parth	MP	X	X							
SV7140CN	50	Yes	Gyn	MP	X	X	X		X	X				
SVCN6404	52	Yes	Gyn	MHP	X	X	X	X	X	X				
Vlaspiik	51	Yes	Gyn	MP	X	X	X		X	X				
Protected Culture / High Tunnels	Corinto	48	Yes	Parth	F	X					X			
	Cucapa	48	Yes	Parth	F		X				X			
	Excelsior	50	Yes	Parth	F	X	X				X			
	Lisboa	60	Yes	Parth	F	X								
	Picolino	45	Yes	Parth	F		X				X			
	Rocky	46	Yes	Parth	F	X	X							
Socrates	52	Yes	Parth	F	X	X								

¹Varieties listed alphabetically within type. ²Hybrid. ³Gyn=Gynoecious or mostly female flowers; 5-15% of a monoecious pollinizer variety added; Mon=Monoecious type with female and male flowers; Parth=Parthenocarpic type that sets fruit without pollination. ⁴F=Fresh Market, P=Processing (pickling), H=Hand harvest multiple times, M=Machine harvest once. ⁵X=high or intermediate level of resistance to Scab, PM=Powdery Mildew, AN=Anthracnose, DM=Downy Mildew, ALS=Angular Leaf Spot, Cmv=Cucumber mosaic virus, Wmv=Watermelon Mosaic Virus, Zmv=Zucchini yellows mosaic virus, Prsv=Papaya ring spot virus. ⁵Only varieties with some resistance to the current strain of downy mildew are noted with an X.

Recommended Nutrients Based on Soil Tests

In addition to using the table below, check the suggestions on rate, timing, and placement of nutrients in your soil test report and chapter B Soil and Nutrient Management. Your state's soil test report recommendations and/or your farm's nutrient management plan supersede recommendations found below.

Cucumbers ¹		Soil Phosphorus Level				Soil Potassium Level				
		Low	Med	High (Opt)	Very High	Low	Med	High (Opt)	Very High	
	N (lb/A)	P ₂ O ₅ (lb/A)				K ₂ O (lb/A)				Nutrient Timing and Method
	80-150	150	100	50	0 ²	200	150	100	0 ²	Total nutrient recommended
	25-50	125	75	25	0 ²	175	125	75	0 ²	Broadcast and disk-in
	25	25	25	25	0	25	25	25	0	Band place with planter
	25-75	0	0	0	0	0	0	0	0	Sidedress when vines begin to run

¹For plasticulture, fertilization rates are based on a standard row spacing of 6 ft. Sulfur at a rate of 20 lb/A in the form of Ammonium Sulfate has been shown to improve color in pickling cucumbers. ²In VA, crop replacement values of 25 lb/A of P₂O₅ and 50 lb/A of K₂O are recommended on soils testing Very High.

Fertigation Schedule Examples

This table provides examples of fertigation schedules based on two common scenarios - sandy coastal plain soils and heavier upland soils. It should be modified according to specific soil tests and base fertility.

Fertigation recommendations for 125 lb N and 125 lb K ₂ O ^{1,2}								
For soils with organic matter content less than 2% or coarse texture and low to medium or deficient K								
			Nitrogen			Potash		
Preplant (lb/A) ³			25			50		
			N	N	N	K ₂ O	K ₂ O	K ₂ O
Stage and Description	Weeks	Days	lb/day	lb/week	lb/stage	lb/day	lb/week	lb/stage
1 Early vegetative	1	1-7	0.5	3.5	3.5	0.4	2.8	2.8
2 Late vegetative	2-3	8-14	0.9	6.3	12.6	0.7	4.9	9.8
3 Fruiting and harvest	4-7	15-42	1.4	9.8	39.2	0.9	6.3	25.2
4 Later harvest ⁴	8-10	43-70	0.9	6.3	18.9	0.6	4.2	12.6

Fertigation recommendations for 75 lb N and 50 lb K ₂ O ^{1,2}								
For soils with organic matter content greater than 2% or fine texture and high or optimum K								
			Nitrogen			Potash		
Preplant (lb/A) ³			50			50		
			N	N	N	K ₂ O	K ₂ O	K ₂ O
Stage and Description	Weeks	Days	lb/day	lb/week	lb/stage	lb/day	lb/week	lb/stage
1 Early vegetative	1	1-7	1	7	7	1	7	7
2 Late vegetative	2-3	8-14	1.5	10.5	21	1.6	11.2	22.4
3 Fruiting and harvest	4-7	15-42	2.2	15.4	61.6	2.2	15.4	61.6
4 Later harvest ⁴	8-10	43-70	1.7	11.9	35.7	1.6	11.2	33.6

¹Rates are based on 7,260 linear bed ft/A (6 ft bed spacing). If beds are closer or wider, fertilizer rates should be adjusted proportionally. Drive rows should not be used in acreage calculations (see section C 3 Fertigation in chapter C Irrigation Management). ²Base overall application rate on soil test recommendations. ³Applied under plastic mulch to effective bed area using modified broadcast method. ⁴For extended harvest after 10 weeks continue fertigation at this rate.

Plant Tissue Testing

Plant tissue testing can be a valuable tool to assess crop nutrient status during the growing season, to aid with in-season fertility programs or to evaluate potential deficiencies or toxicities.

Critical cucumber tissue test values for most recently matured leaves at first bloom are: N 3.5-6 %, P 0.3-0.6 %, K 1.6-3.0 %, Ca 2-4 %, Mg 0.5-0.7% and S 0.3-0.8%. For additional nutrients and other growth stages consult with a tissue testing laboratory or this web link at the University of Florida: <http://edis.ifas.ufl.edu/ep081>

Seed Treatment

Seed should be treated; check with your seed company and see Disease Control below.

Planting Dates

Direct seeding starts late-April in warmer, southern areas and after May 10 in PA and other cool areas. Successive plantings can be made through early August. Container-grown plug plants are started 3 weeks ahead of transplanting. On plastic mulch, planting starts when daily mean temperatures have reached 60°F (16°C). First

F Cucumbers

transplanting dates vary from April 10 in southern regions to June 1 in northern areas. Early plantings should be protected from winds with row covers or rye windbreaks.

Spacing

Slicers: Space rows 3-4 ft apart with plants 9-12 inches apart. Seeding rate: apart with plants 9-12 inches apart for direct seeding bare ground. For plasticulture, space beds 6-8 feet apart and seed or transplant 1- 2 rows per bed, 9-12 inches apart in the row.

Machine Harvest Pickles: Research and field data have shown that 55,000-65,000 plants/A is the optimum population for yield and quality. Parthenocarpic pickles are being trialed in the region. These are planted to achieve 22,000 to 30,000 plants/A.

Hand Harvest Pickles: Space rows 3-4 ft apart with plants 6-8 inches apart. Seeding rate: 1.5-2 lb/A.

Mulching and Fumigation

Plastic mulch laid on moist soil before field planting conserves moisture, and increases soil temperature and early and total yield. Various widths of plastic are available; choose one that works with your production system and equipment. Fumigation will be necessary when there is a history of soil-borne diseases in the field; several fumigants can be used on cucumber depending on what the predominant pests are (see section E 1.5 Soil Fumigation in chapter E Pest Management). Fumigation also aids in the control of weeds. Fumigant and mulch should be applied to well-prepared planting beds; check the fumigant label for the plant-back period that must be adhered to for crop safety. Plastic should be laid immediately over the fumigated soil. Fumigation alone may not provide satisfactory weed control under plastic. Black plastic can be used without a herbicide to provide control of most weeds.

Fertilizer must be applied during bed preparation. At least 50% of the N should be in the nitrate (NO_3^-) form. Drip (trickle) irrigation is recommended for plastic mulch systems and tape is laid at the same time as mulch. Foil and highly reflective mulches can be used to repel aphids that transmit viruses in fall-planted (after July 1) cucurbits. Direct seeding through the mulch is recommended for maximum virus protection; transplants should not be used with foil mulches. Also, an herbicide is not necessary.

Cucumbers also have been successfully grown in no-till systems on cover crop mulch.

Irrigation

Cucumbers require irrigation for best yield and quality. During flowering and fruiting water use can be over 0.25 inches/day and water deficit during this period will have the greatest negative impact on yield and fruit quality. A balance must be struck, however, between maintaining adequate moisture for fruiting while minimizing wetness in the canopy and on the soil surface which promotes fruit rots and downy mildew.

Trellising

Fresh market slicer cucumbers and pickles may be produced on trellises which may result in 2-3 times greater average yield than in non-trellised fields. Trellising is the preferred system in high tunnels. Trellising incurs a higher cost than growing cucumbers on the ground, but it has the following benefits:

1. Improved fruit quality, particularly with respect to color and shape (no yellow "ground spot").
2. More effective control of many diseases and insects.
3. Less damage to vines resulting in a longer harvest season.
4. More consistent and thorough harvesting resulting in fewer jumbos and culls.
5. Easier harvesting than ground grown cucumbers.

Erect the trellis so that it is 6 ft high with a top (No. 8) and bottom (No. 12) wire and plastic twine or netting tied between the two wires at each plant. Posts or poles should be no more than 15 ft apart and the top wire should be very taut. An additional brace between posts may be required when the fruit load becomes heavy. In high tunnels, wires are stretched at the height desired and plastic twine is used to train plants. Training the main stem is required until it reaches and extends over the top wire. Pruning lateral runners near the base of the plant will result in higher yields. The first 4-6 lateral runners that appear should be removed. Other runners above this point should be allowed to run. Single stem systems are often used in high tunnels.

Pollination

Honeybees, squash bees, bumblebees and other wild bees are important for proper cucumber pollination and fruit set. In high tunnels bumblebees are particularly effective. Populations of pollinating insects may be adversely

affected by insecticides applied to flowers or weeds in bloom. Apply insecticides only in the evening hours or wait until bloom is completed before application (see also section A 12 Pollination). Follow all label requirements for pollinator protection. Bee Toxicity ratings are available in the insecticide tables that follow.

Parthenocarpic Cucumbers

Parthenocarpic cucumbers do not require pollination to set fruit. They will be nearly seedless or have unformed seeds. They should be isolated from seeded cucumber types to increase productivity and maintain the seedless nature. Parthenocarpic types should be considered when bee activity is limited such as in high tunnels, under row covers, or in very early plantings.

Season Extension

Low Tunnel Cucumber Production Cucumbers for early production may be successfully grown in high tunnels, in low tunnels with perforated clear plastic row covers, or using floating row covers. Use plastic mulch and trickle irrigation as discussed above. The following field system - similar to that used for early sweet corn - is also successful: A modified bedshaper is used to form a ridge on each side of the plant row, leaving a suitable area for planting. A 36-inch wide piece of embossed clear plastic is then used to cover the plant row, leaving a 5-6 inch high space between the planted row and the plastic cover. It is estimated that temperatures may be increased 10-20°F depending on time of planting and sunlight availability and intensity.

High Tunnel Cucumber Production Cucumbers are a potentially profitable crop for spring and fall production within a high tunnel. Cucumbers mature in approximately half the length of time required for tomato ripening. Cucumbers are also amenable to vertical trellising which increases production and quality. High tunnel cucumber varieties are often parthenocarpic (requiring no pollenizers) although gynocious varieties can also be used (with pollenizers). Cucumbers can be established by direct seeding or transplanting. Space plants 12-18 inches apart in-row on 42-48 inch bed centers. High tunnel varieties can remain unpruned, though pruning can reduce pest infestation and improve marketable yield. If pruning is done, the lower laterals (suckers) should be pruned on the bottom 2 ft leaving 1 or 2 stems per plant to trellis. More information on relative planting and harvesting dates is available in section A 9 High Tunnels in the General Production Recommendations chapter.

Greenhouse Production Varieties are usually parthenocarpic varieties bred specifically for the lower light conditions of fall, winter, and early spring. European “English” or “Dutch” types and Asian types are available. Hydroponic nutrient solution systems are commonly used and cucumbers are trellised with single or double stems trained onto twine; see also section A 10 Greenhouse Production in the General Production Recommendations chapter.

Harvest and Storage

Cucumbers should be harvested when they have reached full size for the variety but while seeds are still soft. For slicers and manually harvested pickles, multiple harvests at 2-3 day intervals will be necessary. Machine-harvested pickles are harvested once when less than 5% have become oversized, as this produces the highest bushel yields. Size requirements of processors will also dictate schedules for machine and hand harvesting pickles.

Cucumbers can be held for 10-14 days at 50-54°F with a relative humidity of 85-90%. At 50°F and above, cucumbers ripen rapidly, with the green color changing to yellow, starting after about 10 days. The color change is accelerated if cucumbers are stored in the same room as apples, tomatoes, or other ethylene-producing crops. Cucumbers for fresh market are usually waxed to reduce moisture loss. Cucumbers are subject to chilling injury if held below 50°F for longer than about 2 days

Weed Control

THE LABEL IS THE LAW-see the Pesticide Use Disclaimer on the first page of chapter F.

Recommended Herbicides

1. Identify the weeds in each field and select recommended herbicides. More information is available in the “Herbicide Effectiveness on Common Weeds in Vegetables” (Table E-2) in chapter E Pest Management.
2. Minimize herbicide resistance development. Identify the herbicide site of action group number and follow recommended good management practices; **bolded group numbers in tables below are herbicides at higher risk for selecting resistant weed populations.** Include non-chemical weed control whenever possible.

F Cucumbers

Labeled Applications Sites for Cucumbers									
		Plastic mulch production					Bare-ground production		
		Soil-Applied		Postemergence			Soil-applied	POST	Post-harvest
Herbicides	WSSA group number	Under Plastic	Row Middles	Over Plastic	Row Middles	Post-Harvest			
Sandea	2	YES	YES	YES	YES		YES	YES	
Treflan	3		YES						
Curbit	3		YES				YES		
Prefar	8	YES	YES				YES		
Command	13		YES				YES		
Strategy	3+13		YES				YES		
Select	1			YES	YES			YES	
SelectMax	1			YES	YES			YES	
Poast	1			YES	YES			YES	
Gramoxone*	22					YES		YES	

*Special Local Needs Label 24(c), be sure it is registered for the specific state and for the intended use.

1. Soil-Applied						
Group	Product Name	Product Rate	Active Ingredient (* = Restricted Use)	Active Ingredient Rate	PHI (d)	REI (h)
2	Sandea 75DF	0.5 to 1 oz/A	halosulfuron	0.023 to 0.047 lb/A	14	12
<p>-Plasticulture: can be applied in a band under the plastic, immediately before laying the mulch; delay seeding or transplanting for 7 days after application. Row row middles: apply before or after weed emergence; apply as a shielded application to avoid contact with the crop. If weeds have emerged, use a non-ionic surfactant at 0.25% v/v or include a non-selective herbicide.</p> <p>-Bareground: apply broadcast after seeding but before crop emergence or no sooner than 7 days before transplanting.</p> <p>-Suppresses or controls yellow nutsedge and certain broadleaf weeds.</p> <p>-Sandea is an ALS inhibiting herbicide and resistant weed populations are common in the region. Do not use Group 2 herbicides repeatedly in the same field. Do not apply Sandea to crops treated with a soil applied organophosphate insecticide, or use a foliar applied organophosphate insecticide within 21 days before or 7 days after a Sandea application.</p> <p>-Maximum Sandea applications per year is 2 and do not exceed 2 oz/A during the crop season.</p>						
3	Curbit 3EC	1 to 3 pt/A	ethalfluralin	0.38 to 1.13 lb/A	--	24
<p>-Plasticulture row middles only: apply as a banded spray after crop emergence or transplanting. Do not soil incorporate.</p> <p>-Bareground: apply broadcast after direct-seeding but prior to crop emergence; do not use on transplanted cucumbers.</p> <p>-Controls annual grasses and certain annual broadleaf weeds, including carpetweed and pigweed sp.</p> <p>-Use lower rate for coarse-textured soils or soils with low organic matter.</p> <p>-Where overhead irrigation is available, activate Curbit with 0.5 inch of irrigation within 2 days after application; if no irrigation or rainfall occurs within 5 days of application, activity of Curbit can be reduced.</p> <p>-Available as a pre-mix herbicide Strategy. Strategy at 3 pt/A= Curbit at 26 fl oz (0.6 lb ai) and Command at 8 fl oz (0.188 lb ai)</p> <p>-Maximum applications per season: not specified</p>						
3	Treflan 4EC	1 to 2 pt/A	trifluralin	0.5 to 1 lb/A	30	12
<p>-Plasticulture row middles only: apply as a directed spray after emergence when plants have reached the 3 to 4 true leaf stage of growth.</p> <p>-Not labeled for bareground production. Primarily controls annual grasses with a few broadleaf weeds. -Do not use (or reduce the rate) when cold, wet soil conditions are expected, or crop injury may result. -Maximum applications per season: not specified.</p>						
3 + 13	Strategy 2.1SC	1.5 to 6 pt/A	ethalfluralin plus clomazone	0.39 to 1.58 lb/A	45	24
<p>-Plasticulture: row middles application. Bareground: apply broadcast just before planting or after planting but before crop emergence.</p> <p>-Strategy is a prepackage mixture of Curbit 3EC and Command 3ME. Refer to individual products for comments.</p> <p>-Clomazone spray or vapor drift may injure susceptible crops and other vegetation, refer to Command 3ME for comments.</p> <p>-Do not apply prior to planting crop. Do not soil incorporate. Maximum applications per season: not specified.</p>						
8	Prefar 4E	5 to 6 qt/A	bensulide	5 to 6 lb/A	45	12
<p>-Plasticulture: under plastic: apply in a band under the plastic, immediately before laying the mulch. Allow 7 day before making transplant holes to allow condensation to incorporate the herbicide. Plasticulture: row middles application is labeled.</p> <p>-Bareground: apply preemergence or preplant incorporated.</p> <p>-Preemergence applications should be followed by irrigation within 36 h (apply enough water to wet the soil at least 2 to 4 inches deep). Preplant incorporated applications should be incorporated 1 to 2 inches deep (deeper than 2 inches will result in reduced weed control).</p> <p>-Provides control/suppression of some annual grass weeds and some broadleaves including pigweeds, purslane, and lambsquarters.</p> <p>-Do not apply more than 6 lb ai/A per season.</p>						
13	Command 3ME	0.4 to 1 pt/A	clomazone	0.015 to 0.375 lb/A	45	12
<p>Plasticulture: row middles application only. -Bareground: apply broadcast just before planting or after planting but before crop emergence. -Supplemental labeling reduces PHI to 30 days (label expires 12/10/2021).</p>						

1. Soil-Applied, Command - continued on next page

1. Soil-Applied, Command - continued

-Use the lower rate when used on coarse-textured soils low in organic matter, when weed pressure is light, or to minimize herbicide carryover that could affect subsequent crops.

-Controls annual grasses and many broadleaf weeds including common lambsquarters, velvetleaf, spurred anoda, and jimsonweed. Carpetweed, morningglory sp., pigweed sp., and yellow nutsedge will not be controlled. Higher rates will improve control (or expand number of species controlled) such as common cocklebur, common ragweed, or jimsonweed (refer to label for specific weeds and rates).

-**WARNINGS:** Command spray or vapor drift may injure sensitive crops and other vegetation up to several hundred yards from the point of application. **Do not** apply adjacent to sensitive crops (see label) or vegetation, or under unfavorable wind or weather conditions. Command may limit subsequent cropping options, see the label. -Available as a pre-mix herbicide Strategy: Strategy at 3 pt/A= Command at 8 fl oz (0.188 lb ai) and Curbit at 26 fl oz (0.6 lb ai) -Maximum number of Command applications per year: 1.

2. Postemergence

Group	Product Name	Product Rate	Active Ingredient (* =Restricted Use)	Active Ingredient Rate	PHI (d)	REI (h)
1	Select 2EC Select Max 0.97EC	6 to 8 fl oz/A 9 to 16 fl oz/A	clethodim	0.094 to 0.13 lb/A	14	24
	Poast 1.5EC	1 to 1.5 pt/A	sethoxydim	0.19 to 0.28 lb/A	3	12

-**Select 2EC:** use crop oil concentrate (COC) at 1% v/v (1 gal/100 gal of spray solution). **Select Max:** use nonionic surfactant (NIS) at 0.25% v/v (1 qt/100 gal of spray solution). **Poast:** Apply with COC at 1.0% v/v. -**The use of COC may increase the risk of crop injury when hot or humid conditions prevail.** To reduce the risk of crop injury, omit additives or switch to NIS when grasses are small and soil moisture is adequate. -Use lower labeled rates for annual grass control and higher labeled rates for perennial grass control.

-Yellow nutsedge, wild onion, wild garlic, and broadleaf weeds will not be controlled.

-Controls many annual and certain perennial grasses, including annual bluegrass, but Poast is preferred for goosegrass control. For best results, treat annual grasses when they are actively growing and before tillers are present. Control may be reduced if grasses are large or under hot or dry weather conditions.

-Repeated applications may be necessary to control certain perennial grasses. If repeat applications are necessary, allow 14 days between applications.

-**Do not** tank-mix with or apply within 2 to 3 days of any other pesticide, unless labeled, as this may increase the risk of crop injury or reduce the control of grasses.

-**Do not** apply more than 8 fl oz of Select 2EC in a single application and **do not** exceed 32 fl oz/A for the season; **do not** apply more than 16 fl oz of Select Max in a single application and **do not** exceed 64 fl oz/A for the season.

-**Do not** apply more than 1.5 pt/A Poast in single application and **do not** exceed 3 pt/A for the season. -Rainfastness is 1 h.

2	Sandea 75DF	0.5 to 1 oz/A	halosulfuron	0.023 to 0.047 lb/A	14	12
---	-------------	---------------	--------------	---------------------	----	----

-**Plasticulture:** broadcast (over the top) or directed to row middles; broadcast for bareground.

-**Bareground:** apply Sandea after the crop has at least 3 to 5 true leaves but before first female flowers appear and no sooner than 14 days after transplanting. If weeds have emerged, use a non-ionic surfactant at 0.25% v/v.

-Suppresses or controls yellow nutsedge and certain broadleaf; control of weeds taller than 3 inches may not be adequate. Sandea will not control common lambsquarters or eastern black nightshade if applied postemergence; for row middle application, tankmix with a non-selective herbicide to increase spectrum of control.

-Sandea provides both residual and postemergence control of susceptible weed species. Effective postemergence control requires an adjuvant. Sandea is an ALS inhibiting herbicide and resistant weed populations are common in the region. **Do not** use Group 2 herbicides repeatedly in the same field.

-**Do not** apply Sandea to crops treated with a soil applied organophosphate insecticide, or use a foliar applied organophosphate insecticide within 21 days before or 7 days after a Sandea application.

-Rainfastness is 4 h. Sandea applications per year is 2 and **do not** exceed 2 oz/A during the crop season

22	Gramoxone 2SL	1.95 pt/A	paraquat*	0.49 lb/A	14	24
----	---------------	-----------	-----------	-----------	----	----

-**A Supplemental Label has been approved for the use of Gramoxone 2SL for postemergence weed control in DE, MD, NJ, PA, and VA.** Row middles as a shielded application. Apply as a directed spray in a minimum of 20 gal spray mix/A to control emerged weeds between the rows after crop establishment. Include a nonionic surfactant at 0.25% v/v.

-Use shields or hoods to prevent spray contact with the crop and low spray pressure (maximum of 30 psi) to reduce small droplets that are prone to drift. See the label for additional information and warnings.

-Rainfastness is 30 min. A maximum of 3 applications per year are allowed.

-**Restricted-use pesticide.** Only certified applicators, who successfully complete the paraquat-specific training, can mix, load or apply paraquat. Application of paraquat "under the direct supervision" of a certified applicator is no longer allowed. Required training link (<http://usparaquattraining.com>); certified applicators must repeat training every three years.

3. Postharvest

Group	Product Name	Product Rate	Active Ingredient (* =Restricted Use)	Active Ingredient Rate	PHI (d)	REI (h)
22	Gramoxone SL 2.0	2.25 to 3 pt/A	paraquat*	0.56 to 0.75 lb/A	--	24

-**A Special Local Needs Label 24(c) has been approved in VA (expires 12/31/2022) and a Supplemental Label in DE for the use of Gramoxone SL 2.0 for postharvest application to desiccate the crop.**

-Apply after the last harvest for bareground or plasticulture. Always include an adjuvant.

3. Postharvest, Gramoxone, continued on next page

F Cucumbers

3. Postharvest, Gramoxone, continued

-Spray coverage is essential for optimum effectiveness. See the label for additional information and warnings.
 -Rainfastness 30 min. A maximum of 2 applications for crop desiccation are allowed.
 -**Restricted-use pesticide.** Only certified applicators, who successfully complete the paraquat-specific training, can mix, load or apply paraquat. Application of paraquat "under the direct supervision" of a certified applicator is no longer allowed. Required training link (<http://usparaquattraining.com>); certified applicators must repeat training every three years.

4. Other Labeled Herbicides These products are labeled but limited local data are available; and/or are labeled but not recommended in our region due to potential crop injury concerns.

Group	Product Name	Active Ingredient (*=Restricted Use)
14	Aim	carfentrazone

Insect Control

THE LABEL IS THE LAW-see the Pesticide Use Disclaimer on the first page of chapter F. Recommended Insecticides

Seed Treatments for Seedcorn Maggot

Control may be achieved by using commercially applied seed treatments containing chlorpyrifos (Lorsban 50W) or thiamethoxam (Farmore DI-400). **Note:** The use of neonicotinoid insecticides (Group 4A) at planting may help reduce seedcorn maggot populations. See also Maggots in section E 3.1 Soil Pests - Detection and Control.

Aphids Note: Aphids transmit multiple viruses.

Apply one of the following formulations:

Group	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
1A	Lannate LV	1.5 to 3.0 pt/A	methomyl* - melon aphid only	1-3	48	H
4A	Neonicotinoid insecticides registered for use on Cucumbers: see table at the end of Insect Control.					
4D	Sivanto Prime or 200SL	21.0 to 28.0 fl oz/A	flupyradifurone - soil	21	4	M
4D	Sivanto Prime or 200SL	7.0 to 14.0 fl oz/A	flupyradifurone - foliar	1	4	M
9B	Fulfill 50WDG	2.75 oz/A	pymetrozine	0	12	L
9B	PQZ	2.4 to 3.2 fl oz/A	pyrifluquinazon	1	12	L
9D	Sefina	3.0 fl oz/A	afidopyropen	0	12	L
21A	Torac	17.0 to 21.0 fl oz/A	tolfenpyrad	1	12	H
28	Exirel	13.5 to 20.5 fl oz/A	cyantraniliprole	1	12	H
28	Harvanta 50SL	10.9 to 16.4 fl oz/A	cyclaniliprole	1	4	H
28	Verimark	6.75 to 13.5 fl oz/A	cyantraniliprole	1	4	H
28 + 6	Minecto Pro	10.0 fl oz/A	cyantraniliprole + abamectin*	7	12	H
29	Beleaf 50SG	2.0 to 2.8 oz/A	flonicamid	0	12	L

Armyworms and Cabbage Loopers

Apply one of the following formulations:

Group	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
1A	Lannate LV	1.5 to 3.0 pt/A	methomyl*	1-3	48	H
3A	Pyrethroid insecticides registered for use on Cucumbers: see table at the end of Insect Control.					
3A + 4A	Endigo ZC	4.0 to 4.5 fl oz/A	lambda-cyhalothrin* + thiamethoxam	1	24	H
5	Entrust SC (OMRI)	4.0 to 8.0 fl oz/A	spinosad	1	4	M
5	Radiant SC	5.0 to 10.0 fl oz/A	spinetoram	1	4	M
11A	Dipel DF, others (OMRI)	0.5 to 2.0 lb/A	<i>Bacillus thuringiensis kurstaki</i>	0	4	N
11A	XenTari (OMRI) (armyworms)	0.5 to 2.0 lb/A	<i>Bacillus thuringiensis aizawai</i>	0	4	N
11A	XenTari (OMRI) (cabbage loopers)	0.5 to 1.0 lb/A	<i>Bacillus thuringiensis aizawai</i>	0	4	N
18	Intrepid 2F	4.0 to 10.0 fl oz/A	methoxyfenozide	3	4	L
22	Avaunt 30WDG, Avaunt eVo	2.5 to 6.0 oz/A	indoxacarb	3	12	H
28	Coragen 1.67SC	3.5 to 7.5 fl oz/A	chlorantraniliprole - soil	1	4	L
28	Coragen 1.67SC	3.5 to 7.5 fl oz/A	chlorantraniliprole - foliar	1	4	L
28	Exirel (armyworms)	7.0 to 13.5 fl oz/A	cyantraniliprole	1	12	H

Armyworms and Cabbage Loopers - continued on next page

Armyworms and Cabbage Loopers - continued

28	Exirel (cabbage loopers)	10.0 to 17.0 fl oz/A	cyantraniliprole	1	12	H
28	Verimark	6.75 to 13.5 fl oz/A	cyantraniliprole	1	4	H
28	Harvanta 50SL	10.9 to 16.4 fl oz/A	cyclaniliprole	1	4	H
28 + 4A	Voliam Flexi	4.0 to 7.0 oz/A	thiamethoxam + chlorantraniliprole	1	12	H
28 + 6	Minecto Pro	5.5 to 10.0 fl oz/A	cyantraniliprole + abamectin*	7	12	H

Cucumber Beetles

Cucumber beetles can transmit bacterial wilt; however, losses from this disease vary greatly between fields and varieties. Pickling cucumbers grown in high-density rows for once-over harvesting can compensate for at least 10% stand losses. On farms with a history of bacterial wilt control adult beetles before they feed extensively on the cotyledons and first true leaves. If foliar insecticides are used, begin spraying shortly after plant emergence and repeat weekly if new beetles continue to invade fields. Treatments may be required until vines begin to run (usually about 3 weeks after plant emergence). Seeds pretreated with a neonicotinoid seed treatment such as Farmore DI-400 should provide up to 14 days of control of cucumber beetle, otherwise, apply one of the following formulations:

Group	Product Name	Product Rate	Active Ingredient(s) (*= Restricted Use)	PHI (d)	REI (h)	Bee TR
1A	Lannate LV	1.5 to 3.0 pt/A	methomyl*	1-3	48	H
1A	Sevin XLR Plus	1.0 qt/A	carbaryl	3	12	H
3A	Pyrethroid insecticides registered for use on Cucumbers: see table at the end of Insect Control.					
4A	Neonicotinoid insecticides registered for use on Cucumbers: see table at the end of Insect Control.					
28	Exirel	20.5 fl oz/A	cyantraniliprole	1	12	H
28	Harvanta 50SL	10.9 to 16.4 fl oz/A	cyclaniliprole	1	4	H

Cutworms See also section E 3.1. Soil Pests - Detection and Control.

Apply one of the following formulations:						
Group	Product Name	Product Rate	Active Ingredient(s) (*= Restricted Use)	PHI (d)	REI (h)	Bee TR
1A	Lannate LV (variegated cutworm)	1.5 pt/A	methomyl*	1	48	H
1A	Lannate LV (granulate cutworm)	1.5 to 3.0 pt/A	methomyl*	1-3	48	H
3A	Pyrethroid insecticides registered for use on Cucumbers: see table at the end of Insect Control.					

Leafminers

Apply one of the following formulations:						
Group	Product Name	Product Rate	Active Ingredient(s) (*= Restricted Use)	PHI (d)	REI (h)	Bee TR
3A	Pyrethroid insecticides registered for use on Cucumbers: see table at the end of Insect Control.					
4A	Neonicotinoid insecticides registered for use on Cucumbers: see table at the end of Insect Control.					
5	Entrust SC (OMRI)	6.0 to 8.0 fl oz/A	spinosad	1	4	M
5	Radiant SC	6.0 to 10.0 fl oz/A	spinetoram	1	4	M
6	Agri-Mek SC	1.75 to 3.5 fl oz/A	abamectin*	7	12	H
17	Trigard 75WSP	2.66 oz/A	cyromazine	0	12	H
28	Coragen 1.67SC	3.5 to 7.5 fl oz/A	chlorantraniliprole - soil	1	4	L
28	Coragen 1.67SC	5.0 to 7.5 fl oz/A	chlorantraniliprole - foliar	1	4	L
28	Exirel	13.5 to 20.5 fl oz/A	cyantraniliprole	1	12	H
28	Verimark	6.75 to 13.5 fl oz/A	cyantraniliprole	1	4	H
28	Harvanta 50SL	10.9 to 16.4 fl oz/A	cyclaniliprole	1	4	H
28 + 6	Minecto Pro	5.5 to 10.0 fl oz/A	cyantraniliprole + abamectin*	7	12	H

Melonworms and Pickleworms

Apply one of the following formulations. When using foliar materials, make one treatment prior to fruit set, and then treat weekly. Check the label for additional instructions when using soil or drip applications.						
Group	Product Name	Product Rate	Active Ingredient(s) (*= Restricted Use)	PHI (d)	REI (h)	Bee TR
1A	Lannate LV	1.5 to 3.0 pt/A	methomyl*	1-3	48	H
1A	Sevin XLR Plus	0.5 to 1.0 qt/A	carbaryl	3	12	H
3A	Pyrethroid insecticides registered for use on Cucumbers: see table at the end of Insect Control.					
3A + 4A	Endigo ZC	4.0 to 4.5 fl oz/A	lambda-cyhalothrin* + thiamethoxam	1	24	H
5	Entrust SC (OMRI)	4.0 to 8.0 fl oz/A	spinosad	1	4	M

Melonworms and Pickleworms - continued on next page

F Cucumbers

Melonworms and Pickleworms - continued

Group	Product Name	Product Rate	Active Ingredient(s)	PHI (d)	REI (h)	Bee TR
5	Radiant SC	5.0 to 10.0 fl oz/A	spinetoram	1	4	M
18	Intrepid 2F	4.0 to 10.0 fl oz/A	methoxyfenozide	3	4	L
22	Avaunt 30WDG, Avaunt eVo	2.5 to 6.0 oz/A	indoxacarb	3	12	H
28	Coragen 1.67SC	3.5 to 7.5 fl oz/A	chlorantraniliprole - soil	1	4	L
28	Coragen 1.67SC	2.0 to 3.5 fl oz/A	chlorantraniliprole - foliar	1	4	L
28	Exirel	7.0 to 13.5 fl oz/A	cyantraniliprole	1	12	H
28	Verimark	5.0 to 10.0 fl oz/A	cyantraniliprole	1	4	H
28	Harvanta 50SL	10.9 to 16.4 fl oz/A	cyclaniliprole	1	4	H
28 + 4A	Durivo	10.0 to 13.0 fl oz/A	thiamethoxam + chlorantraniliprole	30	12	H
28 + 4A	Voliam Flexi	4.0 to 7.0 oz/A	thiamethoxam + chlorantraniliprole	1	12	H
28 + 6	Minecto Pro	5.5 to 10.0 fl oz/A	cyantraniliprole + abamectin*	7	12	H

Mites

Mite infestations generally begin around field margins and grassy areas. **Do not mow or maintain field margins and grassy areas after midsummer since this forces mites into the crop.** Local infestations can be spot-treated. Begin treatment when 10-15% of the crown leaves are infested early in the season, or when 50% of the terminal leaves are infested later in the season. **Note:** Continuous use of carbaryl or a pyrethroid may result in mite outbreaks.

Apply one of the following formulations:						
Group	Product Name	Product Rate	Active Ingredient(s) (* = Restricted Use)	PHI (d)	REI (h)	Bee TR
3A	Pyrethroid insecticides registered for use on Cucumbers: see table at the end of Insect Control.					
6	Agri-Mek SC	1.75 to 3.5 fl oz/A	abamectin*	7	12	H
10B	Zeal Miticide	2.0 to 3.0 oz/A	etoxazole	7	12	L
20B	Kanemite 15SC	31.0 fl oz/A	acequinocyl	1	12	L
21 A	Magister SC	24.0 to 36.0 fl oz/A	fenazaquin	3	12	H
21A	Portal XLO	2.0 pt/A	fenpyroximate	1	12	L
23	Oberon 2SC	7.0 to 8.5 fl oz/A	spiromesifen	7	12	M
28 + 6	Minecto Pro	5.5 to 10.0 fl oz/A	cyantraniliprole + abamectin*	7	12	H
20D	Acramite 50WS	0.75 to 1.0 lb/A	bifenazate	3	12	M

Thrips

Apply one of the following formulations:						
Group	Product Name	Product Rate	Active Ingredient(s) (* = Restricted Use)	PHI (d)	REI (h)	Bee TR
3A	Pyrethroid insecticides registered for use on Cucumbers: see table at the end of Insect Control.					
4A	Neonicotinoid insecticides registered for use on Cucumbers: see table at the end of Insect Control.					
5	Entrust SC (OMRI)	6.0 to 8.0 fl oz/A	spinosad	1	4	M
5	Radiant SC	6.0 to 10.0 fl oz/A	spinetoram	1	4	M
21A	Torac	21.0 fl oz/A	tolfenpyrad	1	12	H
28	Harvanta 50SL	10.9 to 16.4 fl oz/A	cyclaniliprole	1	4	H

Group 3A Pyrethroid Insecticides Registered for Use on Cucumbers

Apply one of the following formulations (check if the product label lists the insect you intend to spray; the label is the law):						
Product Name	Product Rate	Active Ingredient(s) (* = Restricted Use)	PHI (d)	REI (h)	Bee TR	
Asana XL	5.8 to 9.6 fl oz/A	esfenvalerate*	3	12	H	
Baythroid XL	0.8 to 2.8 fl oz/A	beta-cyfluthrin*	0	12	H	
Bifenthrin 2EC, others	2.6 to 6.4 fl oz/A	bifenthrin*	3	12	H	
Danitol 2.4EC	10.67 to 16.0 fl oz/A	fenpropathrin*	7	24	H	
Hero EC	4.0 to 10.3 fl oz/A	zeta-cypermethrin* + bifenthrin*	3	12	H	
Lambda-Cy 1EC, others	2.56 to 3.84 fl oz/A	lambda-cyhalothrin*	1	24	H	
Mustang Maxx	1.28 to 4.0 fl oz/A	zeta-cypermethrin*	1	12	H	
Permethrin 3.2EC, others	4.0 to 8.0 fl oz/A	permethrin*	0	12	H	
Tombstone, others	0.8 to 2.8 fl oz/A	cyfluthrin*	0	12	H	
Warrior II	1.28 to 1.92 fl oz/A	lambda-cyhalothrin*	1	24	H	
Combo products containing a pyrethroid						
Endigo ZC	4.0 to 4.5 fl oz/A	lambda-cyhalothrin* + thiamethoxam (Group 4A)	1	24	H	
Gladiator	19.0 fl oz/A	zeta-cypermethrin* + abamectin* (Group 6)	7	12	H	
Besiege	6.0 to 9.0 fl oz/A	lambda-cyhalothrin* + chlorantraniliprole (Group 28)	1	24	H	

Group 4A Neonicotinoid Insecticides Registered for Use on Cucumbers					
Apply one of the following formulations (check if the product label lists the insect you intend to spray; the label is the law):					
Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
Actara 25WDG	1.5 to 5.5 oz/A	thiamethoxam	0	12	H
Platinum 75SG	1.66 to 3.67 oz/A	thiamethoxam	30	12	H
Admire Pro	7.0 to 10.5 fl oz/A	imidacloprid - soil	21	12	H
Assail 30SG	2.5 to 5.3 oz/A	acetamiprid	0	12	M
Belay 2.13SC	9.0 to 12.0 fl oz/A	clothianidin - soil/drip	21	12	H
Belay 2.13SC	3.0 to 4.0 fl oz/A	clothianidin - foliar (note: PHI: do not make application after 4 th true leaf has unfolded)	see note	12	H
Scorpion 35SL	9.0 to 10.5 fl oz/A	dinotefuran - soil/drip	21	12	H
Scorpion 35SL	2.0 to 7.0 fl oz/A	dinotefuran - foliar	1	12	H
Venom 70SG	5.0 to 7.5 oz/A	dinotefuran - soil/drip	21	12	H
Venom 70SG	1.0 to 4.0 oz/A	dinotefuran - foliar	1	12	H
Combo products containing a neonicotinoid					
Durivo	10.0 to 13.0 fl oz/A	thiamethoxam + chlorantraniliprole (Group 28)	30	12	H
Endigo ZC	4.0 to 4.5 fl oz/A	thiamethoxam + lambda-cyhalothrin* (Group 3A)	1	24	H
Voliam Flexi	4.0 to 7.0 oz/A	thiamethoxam + chlorantraniliprole (Group 28)	1	12	H

Disease Control

THE LABEL IS THE LAW-see the Pesticide Use Disclaimer on the first page of chapter F.
Recommended Fungicides

Nematode Control

See also the chapter E Pest Management (sections E 1.5 Soil Fumigation and E 1.6 Nematode Control), or apply one of the following:

Code	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
1A	Vydate L	0.5 to 1.0 gal/A Incorporate into top 2-4 inches of soil, OR 2.0 to 4.0 pt/A apply 2 w after planting and repeat 2-3 w later.	oxamyl*	1	48	H
7	Velum Prime 4.16SC	6.5 to 6.84 fl oz/A	fluopyram	0	12	--
--	Nimitz 4EC	3.5 to 5.0 pt/A Incorporate or drip-apply 7 d before planting	fluensulfone	n/a	12	N

Seed Treatment Check if seed has been treated with an insecticide and fungicide. If it has not been treated, use a mixture of thiram 480DP (4.5 fl oz/100 lb seed) and an approved commercially available insecticide.

Damping-Off caused by *Pythium*, *Phytophthora*, and *Rhizoctonia*

Code	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
Apply one of the following at-planting (see label for application methods and restrictions):						
Phytophthora and Pythium root rot						
4	Ridomil Gold 4SL	0.5 to 1.0 pt/A	mefenoxam	5	48	N
4	Ultra Flourish 2E	2.0 to 4.0 pt/A	mefenoxam	5	48	N
4	MetaStar 2E AG	4.0 to 8.0 pt/A	metalaxyl	AP	48	N
Phytophthora, Pythium, and Rhizoctonia root rot						
4 + 11	Uniform 3.66SE	0.34 fl oz/1000 ft row. Avoid direct seed contact, which may cause delayed emergence.	mefenoxam + azoxystrobin	AP	0	N
Rhizoctonia root rot						
11	azoxystrobin 2.08F	0.40 to 0.80 fl oz/1000 ft row	azoxystrobin	AP	4	N
Pythium root rot only						
28	Previcur Flex 6F	1.2 pt/A in transplant water, drip irrigation, or direct spray at base of plant and soil	propamocarb HCl	2	12	N

Bacterial and Fungal Diseases

Angular Leaf Spot

Resistant varieties should be used when possible (see table Recommended Varieties). At first sign of disease, apply the labeled rates of fixed copper plus mancozeb. Some coppers are OMRI-approved and can be used in organic production systems to help suppress Angular leaf spot and some fungal diseases. Repeat every 7 days. To minimize the spread of disease, avoid working in field while foliage is wet.

Anthracnose

Resistant varieties should be used when possible (see table Recommended Varieties). Begin fungicide applications when vines begin to run, or earlier if symptoms are detected. Alternate chlorothalonil or mancozeb with other effective fungicides every 7 days. Fungicides with a high risk for resistance development such as FRAC code 11 fungicides that do not come in a mix with another fungicide active ingredient that is effective on anthracnose, should be tank-mixed with a protectant fungicide. Use at least the minimum labeled rate of each fungicide in the tank-mix. **Do not** apply FRAC code 11 fungicides more than 4 times total per season. **Do not** apply FRAC code 11 fungicides if resistance exists in the area; use fungicides with a different FRAC code instead.

Code	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
Under LIGHT or MODERATE disease pressure ALTERNATE:						
M05	chlorothalonil 6F	2.0 to 3.0 pt/A	chlorothalonil	0	12	N
WITH a TANK MIX the following fungicide PLUS mancozeb 75DF 2.0 to 3.0 lb/A OR chlorothalonil 6F 2.0 to 3.0 pt/A:						
1	thiophanate-methyl 70WP	0.5 lb/A	thiophanate-methyl	1	12	N
Under HIGH disease pressure, TANK-MIX one of the following fungicides WITH chlorothalonil 6F 2.0 to 3.0 pt/A:						
3 + 11	Quadris Top 1.67SC	12.0 to 14.0 fl oz/A	difenoconazole + azoxystrobin	0	12	--
7 + 11	Merivon 2.09SC	5.5 fl oz/A	fluxapyroxad + pyraclostrobin	0	12	N
7 + 11	Pristine 38WG	18.5 oz/A	boscalid + pyraclostrobin	0	12	--
11	azoxystrobin 2.08F	11.0 to 15.5 fl oz/A	azoxystrobin	0	4	N
11	Cabrio 20EG	12.0 to 16.0 fl oz/A	pyraclostrobin	0	12	N
AND ROTATE with a TANK-MIX of the following fungicide PLUS mancozeb 75DF 2.0 to 3.0 lb/A OR chlorothalonil 6F 2.0 to 3.0 pt/A every 7 days						
1	thiophanate-methyl 70WP	0.5 lb/A	thiophanate-methyl	1	12	N

Bacterial Wilt

Controlling striped and spotted cucumber beetles is essential for preventing bacterial wilt. See preceding "Cucumber Beetle" section under Insect Control for specific recommendations. Insecticide applications made at seeding may not prevent beetle damage all season; additional foliar insecticide applications may be necessary.

Belly Rot (*Rhizoctonia*)

Code	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
Apply at the 1 to 3 leaf stage. Make a 2nd application 10-14 d later or just prior to vine tip-over (whichever occurs first):						
11	azoxystrobin 2.08F	11.0 to 15.5 fl oz/A	azoxystrobin	1	4	N

Cottony Leak (*Pythium*) - See also Damping off

At planting, apply mefenoxam (Ridomil Gold 4SL, Ultra Flourish 2E) or metalaxyl (MetaStar 2E AG).

Downy Mildew

The pathogen does not overwinter, but introduction to the region can occur early in the year. Newly developed cultivars with resistance or tolerance should be planted where available (see table Recommended Varieties). Even when using resistant cultivars, a good fungicide program is important. However, fungicide efficacy may vary, as strains of the pathogen may vary between seasons.

Scout fields beginning at plant emergence. Strains of downy mildew that infect one cucurbit crop may not affect cucumber. Unnecessary fungicide application can be avoided by not spraying until disease is predicted in the region on cucumber. Begin sprays when vines run or earlier if disease occurrence is predicted for the region (check the Cucurbit Downy Mildew Forecasting website at <http://cdm.ipmpipe.org>). Once the disease has become established in an area, new plantings should receive an application of Ranman, or Previcur Flex at the 1-3 leaf stage. **Preventative applications are much more effective than applications made after disease is detected. In**

addition, spray programs that include fungicides with several different modes of action (FRAC codes) are more effective than programs with few modes of action. For example, alternate Ranman (Code 21) *PLUS* Gavel (Codes M03 + 22), with Orondis Ultra (Codes 49 + 40) *PLUS* chlorothalonil (Code M05). Follow all fungicide label precautions in order to reduce the chance of resistance development.

Code	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
The following are the most effective products. Sprays should be applied on a 7-day schedule.						
Under severe disease conditions spray interval may be reduced IF the label allows.						
ALWAYS tank mix these products with a protectant fungicide (listed below):						
49+40	Orondis Ultra 2.33SC	5.5 to 8.0 fl oz/A	oxathiapiprolin + mandipropamid	0	4	--
49+M05	Orondis Opti 3.37SC	1.75 to 2.5 pt/A	oxathiapiprolin + chlorothalonil	0	12	--
21	Ranman 400SC	2.10 to 2.75 fl oz/A (do not apply with copper; see label)	cyazofamid	0	12	L
28	Previcur Flex 6F	1.2 pt/A	propamocarb HCl	2	12	N
43	Presidio 4SC	4.0 fl oz/A (caution: pathogen is now less sensitive to Presidio)	fluopicolide	2	12	L
M05+22	Zing! 4.9SC	36.0 fl oz/A	chlorothalonil + zoxamide	0	12	N
M05+27	Ariston 42SC	3.0 pt/A	chlorothalonil + cymoxanil	3	12	--
11 + 27	Tanos 50DF	8.0 oz/A	famoxadone + cymoxanil	3	12	--
27	Curzate 60DF	3.2 oz/A	cymoxanil	3	12	N
40	Forum 4.17SC	6.0 fl oz/A	dimethomorph	0	12	N
40 + 45	Zampro 525SC	14.0 fl oz/A	dimethomorph + acetochradin	0	12	--
22	Elumin 4SC	8.0 fl oz/A	ethaboxam	2	12	--
TANK-MIX WITH protectant fungicides:						
M03	mancozeb 75DF	3.0 lb/A	mancozeb	5	24	N
M03+22	Gavel 75DF	1.5 to 2.0 lb/A	mancozeb + zoxamide	5	48	--
M05	chlorothalonil 6F	1.5 to 3.0 pt/A	chlorothalonil	0	12	N

Gummy Stem Blight

Gummy stem blight occurs primarily in the late summer. Fungicides with a high-risk for resistance development such as Pristine (FRAC code 11) should be tank-mixed with a protectant fungicide to reduce the chances for resistance development. Use at least the minimum labeled rate for each fungicide in the tank mix. **Do not** apply FRAC code 11 fungicides more than 4 times total per season. Apply fungicides from a different FRAC code if resistance to FRAC code 11 fungicides exists in the area. Begin sprays when vines begin to run.

Code	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
ALTERNATE one of the following formulations:						
M03	mancozeb 75DF	2.0 to 3.0 lb/A	mancozeb	5	24	N
M05	chlorothalonil 6F	2.0 pt/A	chlorothalonil	0	12	N
WITH A TANK-MIX containing either chlorothalonil or mancozeb PLUS one of the following fungicides:						
3	Proline 480SC	5.7 fl oz/A	prothioconazole	7	12	--
3	tebuconazole 3.6F	8.0 fl oz/A	tebuconazole	7	12	N
3	Rhyme 2.08SC	5.0 to 7.0 fl oz/A	flutriafol	0	12	--
3 + 9	Inspire Super 2.82EW	16.0 to 20.0 fl oz/A	difenoconazole + cyprodinil	7	12	--
3 + 7	Luna Experience 3.34SC	10.0 to 17.0 fl oz/A	tebuconazole + fluopyram	7	12	--
9 + 12	Switch 62.5WG	11.0 to 14.0 oz/A	cyprodinil + fludioxonil	1	12	L
3 + 7	Aprovia Top 1.62EC	10.5 to 13.5 fl oz/A	difenoconazole + benzovindiflupyr	0	12	--
3 + 11	Topguard 4.29SC	5.0 to 8.0 fl oz/A	flutriafol + azoxystrobin	1	12	--
7 + 11	Merivon 2.09SC	5.5 fl oz/A	fluxapyroxad + pyraclostrobin	7	12	N
7 + 11	Pristine 38WG	12.5 to 18.5 oz/A	boscalid + pyraclostrobin	0	12	--
7 + 12	Miravis Prime 3.34SC	9.2 to 11.4 fl oz/A	pydiflumetofen + fludioxonil	1	12	--
11	azoxystrobin 2.08F ¹	11.0 to 15.5 fl oz/A	azoxystrobin	0	4	N
11	Cabrio 20EG ¹	12.0 to 16.0 oz/A	pyraclostrobin	0	12	N

¹azoxystrobin 2.08F and Cabrio 20EG are not recommended in MD, DE and VA due to resistance development.

Phytophthora Crown and Fruit Rot

Different strategies should be used to minimize the occurrence of this disease. Rotate away from susceptible crops (such as cucurbits, peppers, lima and snap beans, eggplants, and tomatoes) for as long as possible, improve field drainage, and apply preplant fumigants. When conditions favor disease development apply fungicides following

F Cucumbers

excellent resistance management practices. Fungicides provide suppression only.

Code	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
Apply one of the following fungicides. Rotate fungicides with different FRAC codes and tank mix with a fixed copper (exception: do not tank mix Ranman 400SC with copper).						
49+40	Orondis Ultra 2.33SC	5.5 to 8.0 fl oz/A	oxathiapiprolin + mandipropamid	0	4	--
49+M05	Orondis Opti 3.37SC	1.75 to 2.5 pt/A	oxathiapiprolin + chlorothalonil	0	12	--
40	Revus 2.08F	8.0 fl oz/A	mandipropamid	0	4	--
40 + 45	Zampro 525SC	14.0 fl oz/A	dimethomorph + acetochradin	0	12	--
22	Elumin 4SC	8.0 fl oz/A	ethaboxam	2	12	--
43	Presidio 4SC ¹	3.0 to 4.0 fl oz/A	fluopicolide	2	12	L
M03+22	Gavel 75DF	1.5 to 2.0 lb/A	mancozeb + zoxamide	5	48	--
M05+22	Zing! 4.9SC	36.0 fl oz/A	chlorothalonil + zoxamide	0	12	N
21	Ranman 400SC	2.75 fl oz/A (plus a non-ionic or organosilicon surfactant; see label for additional precautions)	cyazofamid	0	12	L
40	Forum 4.17SC	6.0 fl oz/A	dimethomorph	0	12	N

¹Presidio may also be applied through the drip irrigation (see supplemental label). Soil drench followed by drip application has given good results in some trials on crown rot caused by *Phytophthora capsici*.

Powdery Mildew

Excellent resistance is available (see table Recommended Varieties). The fungus that causes cucurbit powdery mildew has developed resistance to high-risk fungicides. In the Eastern US, resistance to strobilurin (FRAC code 11), DMI (FRAC code 3), and SDHI (FRAC code 7) fungicides has been reported. Proper fungicide resistance management should be followed to help delay the development of resistance and minimize control failures.

Powdery mildew generally occurs from mid-July until the end of the season. Observe plants for the presence of powdery mildew. If one lesion is found on the underside of 45 old leaves/A, begin the following fungicide program:

Code	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
TANK MIX one of these products with a protectant such as chlorothalonil 6F at 2.0 to 3.0 pt/A:						
50	Vivando 2.5SC	15.4 fl oz/A	metrafenone	0	12	--
3 + 7	Luna Experience 3.34SC	10.0 to 17.0 fl oz/A	tebuconazole + fluopyram	7	12	--
AND ALTERNATE with a TANK MIX of one of the following with a protectant such as chlorothalonil 6F at 2.0 to 3.0 pt/A						
3	Proline 480SC	5.7 fl oz/A	prothioconazole	7	12	--
3	Procure 480SC	4.0 to 8.0 fl oz/A	triflumizole	0	12	N
3	Rally 40WSP	5.0 oz/A PLUS	myclobutanil	0	24	N
3	tebuconazole 3.6F	4.0 to 6.0 fl oz/A	tebuconazole	7	12	N
3	Rhyme 2.08SC	5.0 to 7.0 fl oz/A	flutriafol	0	12	--
3 + 9	Inspire Super 2.82EW	16.0 to 20.0 fl oz/A	difenoconazole + cyprodinil	7	12	--
3 + 7	Aprovia Top 1.62EC	10.5 to 13.5 fl oz/A	difenoconazole + benzovindiflupyr	0	12	--
3 + 11	Topguard 4.29SC	5.0 to 8.0 fl oz/A	flutriafol + azoxystrobin	1	12	--
7 + 11	Pristine 38WG	12.5 to 18.5 oz/A	boscalid + pyraclostrobin	0	12	--
U13	Gatten 5EC	6.0 to 8.0 fl oz/A	flutianil	0	12	--
P05	Regalia (OMRI)	4.0 qt/A	Extract of <i>Reynoutria sachalinensis</i>	0	4	--
39	Magister 1.6SC	24 to 36 fl oz/A	fenazaquin	3	12	H
7 + 12	Miravis Prime 3.34SC	9.2 to 11.4 fl oz/A	pydiflumetofen + fludioxonil	1	12	--
U06	Torino 0.85SC	3.4 fl oz/A	cyflufenamid	0	4	--

Scab Scab typically occurs during cool periods. Excellent resistance is available in some varieties and they should be used when possible.

Code	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
Apply one of the following as true leaves form and repeat every 5-7 days:						
M03	mancozeb 75DF	2.0 to 3.0 lb/A	mancozeb	5	24	N
M05	chlorothalonil 6F	2.0 to 3.0 pt/A	chlorothalonil	0	12	N

Viruses The most prevalent virus in the mid-Atlantic region is WMV2, followed by PRSV, ZYMV and CMV. Use varieties with multiple virus resistance when possible (see table Recommended Varieties). Plant fields far away from existing cucurbit plantings to help reduce aphid transmission of viruses into new fields.

For Immediate Medical Attention

Call 911

**For a Pesticide Exposure Poisoning
Emergency Call**



For All States

This number will automatically connect you to the poison center nearest to you.

Anyone with a poisoning emergency can call the toll-free telephone number for help. Personnel at the Center will give you first-aid information and direct you to local treatment centers if necessary.

For Pesticide Spills

Small Spills: See the product label for cleanup advice.

Large spills: Call the National Response Center at 1-800-424-8802 or CHEMTREC at 800-424-9300 (24 hours) - Industry assistance with emergency response cleanup procedures for large, dangerous spills.

Be aware of your responsibility to report spills to the proper state agency.