

## F. Commodity Recommendations

### Pesticide Use Disclaimer

#### **THE LABEL IS THE LAW**

**Before using a pesticide, check the label for up to date rates and restrictions.**

**Labels can be downloaded from:** <http://www.cdms.net/>, <https://www.greenbook.net/> or <http://www.agrian.com/labelcenter/results.cfm>

**For more information on Pesticide Safety and the Pesticide Label see chapter D.**

#### **Guide to the Recommended Pesticide Tables in the Following Crop Sections:**

**1. Pesticides are listed by group or code number based on chemical structure and mechanism of action**, as classified by the Weed Science Society of America (WSSA) for herbicides, the Insecticide Resistance Action Committee (IRAC) for insecticides, and the Fungicide Resistance Action Committee (FRAC) for fungicides.

**If the number is in bold font, the product may have resistance concerns.**

**2. For restricted use pesticides**, the restricted active ingredients are labeled with a \*. (See section D 3.2.1 “Restricted Use Classification Statement” for more information).

**3. In addition to the pesticides listed below, other formulations or brands with the same active ingredient(s) may be available. ALWAYS CHECK THE LABEL:**

- a) to ensure a pesticide is labeled for the same use,**
- b) to ensure the pesticide is labeled for the desired crop, and**
- c) for additional restrictions.**

**4. All pesticide recommendations are made for spraying a broadcast area of 1 acre** (43,560 square feet). **Adjust the rate for banded applications** (for more information, see section E 1.3 Calibrating Granular Applicators).

**5. Check the label for the maximum amount of pesticide per application and the maximum number of applications per year.**

**6. Bee Toxicity Rating (Bee TR):** N=nontoxic; L=minimum impact on bees; M=moderately toxic, can be used if dosage, timing and method of application are correct, but should NOT be applied directly to the crop if bees are present; H=highly toxic, severe losses expected, -- = data not available.

# Tomatoes

## Recommended Varieties

Listed in alphabetical order within type

### Market Tomatoes<sup>1</sup>

Type	Variety	Color	Season	Culture	Use <sup>2</sup>	Disease Resistance <sup>3</sup>	Plant Habit <sup>4</sup>
Globe	Amelia	Red	Mid	Field	LW, S	V,F,Tswv	D
	BHN 589	Red	Mid	Field, High Tunnel	DM, LW	V,F,Tomv	D
	BHN 602	Red	Mid, Late	Field	DM, LW, S	V,F,Tswv	D
	BHN 871	Yellow	Mid	Field, High Tunnel	DM, LW	V,F,Tomv	D
	BHN 964	Red	Mid	Field	DM, LW, S	V,F,Tomv,Eb	D
	Biltmore	Red	Mid	Field	DM, LW,	V,F,Asc,Gls	D
	Camaro	Red	Mid	Field	LW, S	V,F,Gls, Asc, Tylc	D
	Carolina Gold	Yellow	Mid	Field	DM, LW	V,F	D
	Charger	Red	Mid	Field, High Tunnel	DM, LW, S	V,F,Gls,Asc,Tylc	D
	Defiant	Red	Mid	Field	DM, LW	V,F,Lb, Eb	D
	Dixie Red	Red	Mid	Field	DM, LW, S	V,F,N,Gls,Tswv,Asc	D
	Florida 47R	Red	Mid	Field	LW, S	V,F,Asc,Gls	D
	Florida 91	Red	Mid, Late	Field	DM, LW, S	V,F,Asc,Gls	D
	Grand Marshall	Red	Mid	Field, High Tunnel	DM, LW, S	V,F,Gls, Asc, Tylc	D
	Lemon Boy	Yellow	Mid	Field, High Tunnel	DM, LW	V,F,N	I
	Mountain Fresh Plus	Red	Mid, Late	Field	DM, LW, S	V,F,N	D
	Mountain Glory	Red	Mid	Field	DM, LW, S	V,F,Gls,Tswv	D
	Mountain Merit	Red	Mid	Field	DM, LW, S	V,F,N,Tswv, Lb,	D
	Mountain Spring	Red	Mid	Field	DM, LW	V,F	D
	Phoenix	Red	Mid, Late	Field	LW, S	V,F,Asc,Gls	D
	Primo Red	Red	Early	Field	DM, LW, S	V,F,Tomv	D
	Red Bounty	Red	Mid, Late	Field, High Tunnel	DM, LW	V,F,N,Gls,Tswv	D
	Red Defender	Red	Mid	Field	DM, LW, S	V,F,N,Tswv	D
	Red Deuce	Red	Mid	Field	DM, LW, S	V,F,Tomv,Gls,Asc	D
	Red Morning	Red	Mid	Field	DM, LW, S	V,F, Tomv, Tswv	D
	Red Mountain	Red	Mid	Field, High Tunnel	DM, LW, S	V,F,Tswv	D
	Rocky Top	Red	Mid	Field, High Tunnel	DM, LW, S	V,F,Gls	D
	Scarlet Red	Red	Mid	Field, High Tunnel	DM, LW, S	V,F	D
Sunbrite	Red	Early	Field, High Tunnel	DM, LW, S	Asc, V,F,Gls	D	
Volante	Red	Mid	Field	DM, LW, S	V,F,Gls,Asc, Tswv	D	

<sup>1</sup>All varieties are hybrids. <sup>2</sup>DM=Direct Market, LW=Local Wholesale, S=Shipping. <sup>3</sup>Resistances or tolerances: Asc=Alternaria stem canker, Eb=Early blight, F=Fusarium wilt, Gls=Gray leaf spot, Lb=Late blight, N=Root-knot nematode, Tomv=Tomato mosaic virus, Tswv=Tomato spotted wilt virus, Tylc=Tomato Yellow Leaf Curl virus, V=Verticillium wilt. <sup>4</sup>D=Determinate, I=Indeterminate.

### Heirloom Tomatoes

Type	Variety	Color	Size	Maturity	Plant Habit
Beefsteak	Brandywine Red	Red skin, red flesh	Large	Late	I, potato leaf
	Mortgage Lifter	Pink skin, Pink flesh	Large	Late	I
Globe	Cherokee Purple	Burgundy	Medium-Large	Mid	I
	Prudens Purple	Deep pink skin and flesh	Large	Mid	I, potato leaf
Round	Eva Purple Ball	Deep pink skin and flesh	Medium	Mid	I
	Green Zebra	Yellow-gold with dark green strips	Medium	Mid	I
Small pear	Yellow Pear	Yellow skin and flesh	Small	Late	I

### Cherry, Grape, Plum and Cluster Tomatoes

Type	Variety	Color	Disease Resistance <sup>2</sup>	Plant Habit <sup>3</sup>
Cherry	BHN 762	Red	V,F	D
	Sun Gold	Orange	F, Tomv	I
	Sun Sugar	Orange	F, Tmv	I
	Sweet Chelsea	Red	V,F,N,Tomv	I
	Sweet Treats	Pink	F,Tomv,Gls	I
Grape	BHN 784	Red	F	D
	Cupid	Red	F, Asc	I
	Jolly Girl	Red	V, F	D
	Mini Charm	Red	V,F,Tomv	I
	Smarty	Red	V, F	I
	Valentine <sup>4</sup>	Red	Eb	I
Large Grape	Juliet	Red	Eb, Lb	I
Plum	Daytona	Red	Asc, F, N, V	D
	Health Kick	Red	V,F,Asc,Tswv,Bs	D
	Mariana	Red	V,F,N,Asc	D
	Picus	Red	V,F,Asc,Gls,Tswv	D
	Plum Crimson	Red	V,F	D
	Plum Dandy	Red	V,F	D
	Plum Regal	Red	V,F,Lb,Tswv,	D
	Pony Express	Red	V,F,N,Tomv,Bs	D
Small cluster	Mt. Magic	Red	V,F,Lb	I

<sup>1</sup>All varieties are hybrids. <sup>2</sup>Resistances or tolerances: Asc=Alternaria stem canker, Bs=Bacterial speck, Eb=Early blight, F=Fusarium wilt, GlS=Gray leaf spot, Lb=Late blight, N=Root-knot nematode, Tmv=Tobacco mosaic virus, Tomv=Tomato mosaic virus, Tswv=Tomato spotted wilt virus, V=Verticillium wilt. <sup>3</sup>D=Determinate, I=Indeterminate. <sup>4</sup>High lycopene.

### Processing Tomatoes

Type	Variety	Season	Disease Resistance
Processing	TSH4	Early	V,F,Bs
	H-3402	Mid	V,F,N,Bs
	H-3406	Full	V,F,Bs,Eb,Bc
	H-9997	Early	V,F,N,Asc,Bs

All processing varieties are hybrids. Most plantings are contracted by processor; consult with processor to determine preferred varieties. Disease resistance or tolerance: Asc=Alternaria stem canker, Bs = Bacterial speck, Bc=Bacterial Canker, Eb = Early Blight F=Fusarium wilt, N=Root-knot nematode, V=Verticillium wilt.

### Recommended Nutrients Based on Soil Tests

In addition to using the table below, check the suggestions on rate, timing, and placement of nutrients in your soil test report and chapter B Soil and Nutrient Management. Your state’s soil test report recommendations and/or your farm’s nutrient management plan supersede recommendations found below.

Tomatoes <sup>1</sup>	N (lb/A)	Soil Phosphorus Level				Soil Potassium Level				Nutrient Timing and Method
		Low	Med	High (Opt)	Very High	Low	Med	High (Opt)	Very High	
		P <sub>2</sub> O <sub>5</sub> (lb/A)				K <sub>2</sub> O (lb/A)				
Bare-Ground Fresh Market	80-90	200	150	100	0 <sup>2</sup>	300	200	100	0 <sup>2</sup>	Total nutrient recommended
	40-45	200	150	100	0 <sup>2</sup>	300	200	100	0 <sup>2</sup>	Broadcast and disk-in
	40-45	0	0	0	0	0	0	0	0	Sidedress when first fruits are set
Processing Machine Harvest	50-75	200	150	100	0 <sup>2</sup>	250	150	100	0 <sup>2</sup>	Total nutrient recommended
	25	200	150	100	0 <sup>2</sup>	250	150	100	0 <sup>2</sup>	Broadcast and disk-in
	25-50	0	0	0	0	0	0	0	0	Sidedress at first cultivation
Polyethylene Mulched Fresh Market	150-210	200	150	100	0 <sup>2</sup>	300	200	100	0 <sup>2</sup>	Total nutrient recommended
	0	200	150	100	0 <sup>2</sup>	150	100	50	0	Broadcast and disk-in
	50-85	0	0	0	0	0	0	0	0	Incorporate into the plant bed before laying polyethylene mulch
	90-125	0	0	0	0	150	100	50	0 <sup>2</sup>	Fertigate 0.5 to 2.5 lb/day. See chart and Drip/Trickle Fertilization section

<sup>1</sup>Apply 1-2 lb/A of boron (B) with broadcast fertilizer; see also Table B-7 in chapter B Soil and Nutrient Management.

<sup>2</sup>In VA, crop replacement values of 50 lb/A of P<sub>2</sub>O<sub>5</sub> and 50 lb/A of K<sub>2</sub>O are recommended on soils testing Very High.

## Drip/Trickle Fertigation

Before laying plastic mulch, adjust soil pH to 6.5 and broadcast and disk in preplant nutrients (see table above) Apply the balance of your needed K<sub>2</sub>O that you do not plan to apply via fertigation as a modified broadcast application that treats only the mulched area. Nitrogen fertilizer should be incorporated into the bed or split between incorporated and a surface band bed treatment immediately before laying plastic mulch.

After laying plastic mulch and installing the trickle irrigation system, apply completely soluble fertilizer through the drip system to supply additional N and potash throughout the season and adjust rates as necessary based on soil and tissue tests (see tables below). For more information, see **Fertigation Rates for Drip Irrigated Plasticulture Crops** in chapter C Irrigation Management, section C 3 Fertigation.

## Fertigation Schedule Examples for Fresh Market Tomatoes

This table provides examples of fertigation schedules based on two common scenarios – sandy coastal plain soils and heavier upland soils. Modify according to specific soil tests and base fertility.

Fertigation recommendations for 150 lb N and 150 lb K <sub>2</sub> O <sup>1,2</sup>								
For soils with organic matter content less than 2% or coarse texture and low to medium or deficient K								
			Nitrogen			Potash		
Preplant (lb/A) <sup>3</sup>			50			125		
			N	N	N	K <sub>2</sub> O	K <sub>2</sub> O	K <sub>2</sub> O
Stage and Description	Weeks	Days	lb/day	lb/week	lb/stage	lb/day	lb/week	lb/stage
1 Early vegetative	1-2	1-14	0.5	3.5	7	0.5	3.5	7
2 Late vegetative	3-4	15-28	0.7	4.9	9.8	0.7	4.9	9.8
3 Early flowering	5-6	29-42	1.0	7	14	1	7	14
4 Flowering and fruiting	7-8	43-56	1.5	10.5	21	1.5	10.5	21
5 Early harvest	9-11	57-77	2.2	15.4	46.2	2.2	15.4	46.2
6 Later harvest <sup>4</sup>	12-14	78-98	2.5	17.5	52.5	2.5	17.5	52.5

  

Fertigation recommendations for 75 lb N and 75 lb K <sub>2</sub> O <sup>1,2</sup>								
For soils with organic matter content greater than 2% or fine texture and high or optimum K								
			Nitrogen			Potash		
Preplant (lb/A) <sup>3</sup>			50			50		
			N	N	N	K <sub>2</sub> O	K <sub>2</sub> O	K <sub>2</sub> O
Stage and Description	Weeks	Days	lb/day	lb/week	lb/stage	lb/day	lb/week	lb/stage
1 Early vegetative	1-2	1-14	0.25	1.75	3.5	0.25	1.75	3.5
2 Late vegetative	3-4	15-28	0.35	2.45	4.9	0.35	2.45	4.9
3 Early flowering	5-6	29-42	0.5	3.5	7	0.5	3.5	7
4 Flowering and fruiting	7-8	43-56	0.75	5.25	10.5	0.75	5.25	10.5
5 Early harvest	9-11	57-77	1.1	7.7	23.1	1.1	7.7	23.1
6 Later harvest <sup>4</sup>	12-14	78-98	1.25	8.75	26.25	1.25	8.75	26.25

<sup>1</sup>Rates above are based on 7,260 linear bed ft/A (6 ft bed spacing). If beds are closer or wider, fertilizer rates should be adjusted proportionally. Drive rows should not be used in acreage calculations. See section C 3 Fertigation in the Irrigation Management chapter for more information.

<sup>2</sup>Base overall application rate on soil test recommendations. <sup>3</sup>Applied under plastic mulch to effective bed area using modified broadcast method. <sup>4</sup>For extended harvest after 10 weeks continue fertigation at this rate.

## Plant Tissue Testing

Plant tissue testing can be a valuable tool to assess crop nutrient status during the growing season to aid with in-season fertility programs or to evaluate potential deficiencies or toxicities. Critical tomato tissue test values for most recently matured leaves at first flower are: 2.8-4.0 %, P 0.2-0.4 %, K 2.5-4.0 %, Ca 1.0-2.0 %, Mg 0.25-0.5% and S 0.3-0.6%. For additional nutrients and other growth stages consult with a tissue testing laboratory or this web link at the University of Florida: <http://edis.ifas.ufl.edu/ep081>.

## Plant Petiole Sap Testing

Plant petiole sap and tissue testing are valuable tools to assess crop nutrient status during the growing season, to aid with in-season fertility programs, or to evaluate potential deficiencies or toxicities.

Tomato Developmental Stage	Fresh Petiole Sap Concentration (ppm)	
	NO <sub>3</sub> -N	K
First buds	1000-1200	3500-4000
First open flowers	600-800	3500-4000

*Plant Petiole Sap Testing - continued on next page*

## F Tomatoes

### Plant Petiole Sap Testing - continued

Tomato Developmental Stage	Fresh Petiole Sap Concentration (ppm)	
	NO <sub>3</sub> -N	K
Fruits 1 inch diameter	400-600	3000-3500
Fruits 2 inch diameter	400-600	3000-3500
First harvest	300-400	2500-3000
Second harvest	200-400	2000-2500

## Seed Treatment

Purchase hot water treated seed if possible or request hot water seed treatment. Hot water treatment is administered to eradicate bacterial pathogens. For more information see Disease Control below.

## Grafting

Grafting of fresh-market tomatoes into rootstocks that confer resistance or tolerance to soil-borne pests and diseases is practiced as an alternative to chemical soil fumigation and may increase plant vigor impacting water and nutrient requirements, plant density, pruning, and training, as well as fruit yield and quality. Grafting tomato plants into vigorous rootstocks is also used to enhance plant tolerance to abiotic stress conditions including non-optimal temperatures, excess or deficiency of nutrients, salinity, drought, or alkalinity stress.

### Rootstocks commonly used for tomato, and their disease resistance or susceptibility<sup>1</sup>

Major Tomato Disease	Tomato Mosaic Virus	Fusarium Crown Rot ( <i>Fusarium oxysporum</i> f.sp. <i>radicilycopersici</i> )	Fusarium Wilt ( <i>Fusarium oxysporum</i> f. sp. <i>lycopersici</i> ) <b>Race 1</b>	Fusarium Wilt ( <i>Fusarium oxysporum</i> f. sp. <i>lycopersici</i> ) <b>Race 2</b>	Verticillium Wilt ( <i>Verticillium albo-atrum</i> & <i>V. dahliae</i> )	Bacterial Wilt ( <i>Ralstonia solanacearum</i> )	Root-knot Nematodes ( <i>Meloidogyne</i> spp.)
Anchor-T	R	N/A	R	R	R	R	R
Arnold	R	R	R	R	N/A	N/A	R
Beaufort	R	R	R	R	R	S	R
Body	R	N/A	S	R	R	S	R
Estamino	R	R	R	R	R	S	R
Maxifort	R	R	R	R	R	S	R
Multifort	S	S	R	R	S	S	R
RT-04-105T	S	R	R	R	R	R	R
RT-04-106T	R	S	R	R	S	R	R

<sup>1</sup>R = Resistant, S = Susceptible, N/A-not characterized.

## Hardening Transplants

Hardening seedlings before field planting is recommended. However, hardening by exposure to cool temperatures 60-65°F (16-18°C) day and 50-60°F (10-16°C) night for one week or more causes catfacing. Instead, harden plants by withholding N and reducing water; allow plants to wilt slightly between light waterings.

## Fresh Market

Yield and fruit size and quality of fresh market tomatoes are increased by the use of black plastic mulch in combination with trickle irrigation. Form raised, dome-shaped beds to aid in disease control. Lay 4 ft wide black plastic mulch tightly over the beds. For early summer harvest of market tomatoes, start transplanting April 10-20 in southern or normally warmer areas, and May 10-25 in cooler, northern areas.

**Ground Culture:** Space determinate vined varieties in rows 4-5 ft apart with plants 15-24 inches apart in the row. For indeterminate varieties, space rows 5-6 ft apart with plants 24-36 inches apart in the row.

**Stake Culture:** Staking tomatoes is a highly specialized production system. Staking improves fruit quality by keeping plants and fruit off the ground and allows for better spray coverage. Staked tomatoes are easier to harvest than non-staked tomatoes. The recommendations below are for the short-stake cultural system using determinate cultivars that grow 3-4 ft tall. Row widths of 5-6 ft with in-row spacings of 18-24 inches between plants are recommended.

**Pruning** is practiced to establish a desired balance between vine growth and fruit growth. Little to no pruning results in a plant with a heavy load of smaller fruit. Moderate pruning results in fewer fruit that are larger and easier to harvest. Pruning can result in earlier maturity of the crown fruit and improve spray coverage and pest control.

The pruning method is variety and fertility dependent. Less vigorous determinate cultivars generally require less pruning. Growers should experiment with several degrees of pruning on a small scale to determine pruning requirements for specific cultivars and cultural practices.

Removing all suckers up to the one immediately below the first flower cluster is adequate for most determinate cultivars. Removing the sucker immediately below the first flower cluster or pruning above the first flower cluster can result in severe leaf curling and stunting of the plant. Prune when the suckers are 2-4 inches long. A 2<sup>nd</sup> pruning may be required to remove suckers that are too small to be easily removed during the 1<sup>st</sup> pruning and to remove ground suckers that may develop. Pruning when suckers are too large requires more time and can damage the plants, delay maturity, and increase disease incidence. Do not prune plants when they are wet to avoid spread of bacterial diseases. Pruning should be done before the first stringing because the string can slow down the pruning process.

**Staking** involves setting up a series of wooden stakes with twine woven around the stakes to train the plants to grow vertically off the ground. Stakes 4-4½-ft long by 1-inch square are driven approximately 12 inches into the soil between the plants.

Vigorous cultivars may require larger and longer stakes. A stake placed between every other plant is adequate to support most determinate varieties. Placing an additional stake at an angle and tied to the end stake of each section or row is needed to strengthen the trellis system. Stakes can be driven by hand with a homemade driving tool or with a commercially available, power-driven stake driving tool. Drive stakes to a consistent depth so that spray booms can be operated in the field without damaging the trellis system. Select "tomato twine" that is resistant to weathering and stretching and that binds well to the wooden stakes. Tomato twine is available in 3-4-lb boxes and approximately 30 lb/A are required. To make tying convenient, use a homemade stringing tool made from a length of metal conduit, PVC pipe, broom handle, or wooden dowel. With conduit or PVC pipe, the string is fed through the pipe. With a broom handle or wooden dowel, two small parallel holes, each approximately ½-1 inch from the end, must be drilled to feed the string through one hole along the length of the tool and through the other hole. The tool serves as an extension of the worker's arm (the length cut to the worker's preference) and helps to keep the string tight.

**Stringing** consists of tying the twine to an end stake passing the string along one side of the plants, looping the twine around each stake until you reach the end of a row or section (100-ft sections with alleys may be helpful for harvesting). The same process is continued on the other side of the row. The string tension must be tight enough to hold the plants upright but harvest can be difficult and strings can scar fruit if they are too tight.

The first string should be strung 8-10 inches above the ground when plants are 12-15 inches tall and before they fall over. Run the next string 6-8 inches above the preceding string before plants start to fall over. Three to 4 stringings are required for most determinate varieties. Stringing should be done when the foliage is dry to prevent the spread of bacterial diseases.

## Processing Tomatoes

**Transplanting:** Processing tomatoes can be transplanted starting April 15-20 in warmer, southern areas to May 5-10 in PA and normally cooler areas. Successive plantings can be made through early June. Space transplants 9-12 inches apart in single rows 5 ft. apart or to accommodate machine harvesters. Small, determinate varieties may be grown in double rows. Space double rows 12 inches apart and space plants 12-18 inches apart in each of the double rows.

**Fruit Ripening:** Ethephon is a growth regulator labeled for use on processing tomatoes. Proper application increases earliness and yield and decreases sorting of green fruit in machine-harvested tomatoes. Rate and time of application are critical for successful use, see state fact sheets and check product label for details.

## Harvest and Post-Harvest Considerations

Depending on marketing requirement, tomatoes may be harvested at the **mature green stage** (when and after which the fruit cavity is filled by gel), **breaker stage** (just showing pink at the bottom of the fruit), **semi-ripe** (with different amounts of red pigmentation) or **fully ripe**. Fruit are very perishable and subject to surface and internal damage and must be handled with care. Tomatofields should be harvested often and thoroughly to hasten the ripening of later fruits and reduce the range of ripeness if a specific stage is desired. Harvesting every day may be required during peak season. Remove all diseased, misshapen, and otherwise cull tomatoes from the vines as soon as they are discovered. Remove discarded tomatoes from the field to avoid the spread and buildup of diseases and insect pests. For standard slicing tomatoes, cherry tomatoes, and plum tomatoes, remove the stem during picking. Cluster tomatoes are harvested with the whole truss attached to fruits.

## F Tomatoes

Tomatoes should be washed sufficiently to remove dust and foreign material, by hand or mechanically by spraying them with chlorinated water as they move over a set of soft brush rolls. The small amount of retained water may be removed by absorbent rollers alone or in combination with an overhead air-blast drier. The wash water should be several degrees warmer than the pulp temperature of the tomatoes to avoid drawing water and disease organisms into the fruit. The water should be chlorinated at the rate of 125 ppm. The chlorine level and pH (6 to 7) of the wash water should be checked at least hourly during the day with test papers or a meter. Tomatoes are then sized and separated by color and grade and carefully packed into 25 lb boxes.

### Size Classification of Tomatoes

Size Designation	Minimum Diameter (inch)	Maximum Diameter (inch)
Extra small	1-28/32	2-4/32
Small	2-4/32	2-9/32
Medium	2-9/32	2-17/32
Large	2-17/32	2-28/32
Extra large	2-28/32	3-15/32

### Color Classification of Tomatoes

Tomatoes may be graded into the following color classes (some classes may be combined).

<b>Green</b>	The surface of the tomato is completely green. The shade of green may vary from light to dark. Mature green fruits are typically ripened at the terminal market or by the repacker with ethylene gas.
<b>Breakers</b>	There is a definite break in the color from green to tannish yellow with pink or red skin covering not more than 10% of the surface.
<b>Turning</b>	More than 10% but not more than 30% of the surface, shows a definite change in color from green to tannish yellow, pink, red, or a combination of those colors.
<b>Pink</b>	More than 30% but not more than 60% of the surface shows pinkish red or red color.
<b>Light Red</b>	More than 60% but not more than 90% shows pinkish red or red color.
<b>Red</b>	More than 90 % of the surface shows red color.

### Shipping

For long distance shipping, mature green harvest is the common practice. For local wholesale, harvest is usually at the breaker stage. For direct market, harvest is at the ripe stage. Store mature-green tomatoes at 55-70°F (13-21°C); breakers, partially ripe, and ripe fruit at 50°F (10°C) and a relative humidity of 90-95%. Exposing tomatoes to temperatures below 50°F results in loss of color, shelf life, firmness and flavor.

### Tomato Disorders

#### **Blossom-End Rot (BER)**

This physiological disorder is caused by inadequate movement of calcium into the fruit. BER occurs at low soil moisture and is more severe when plants have small, shallow root systems. Plastic mulch can restrict the movement of water to the root zone and increase BER. Hot, windy conditions increase water loss from the plant and increase the incidence of BER.

Be sure soil calcium is sufficient and in balance with other essential plant nutrients. Test the soil and apply calcitic lime and fertilizer according to recommendations, then lay plastic mulch when soil moisture is optimal for planting. Apply irrigation to wet the root zone and encourage deep root development.

#### **Blotchy Ripening, Graywall and Internal White Tissue**

These problems are a complex of physiological disorders and pathological diseases. Blotchy Ripening and graywall often appear on shaded fruit growing in the interior of dense vegetative plants. Yellow-eye, a ring of yellow tissue surrounding the blossom scar, often occurs in fruit with blotchy ripening and internal white tissue.

*Blotchy ripening* is when areas of the fruit do not ripen or do so after the rest of the fruit is ripe. White or yellow blotches may appear on the surface of the fruit while the internal tissue is still hard. Usually this disorder occurs on the upper portion of the fruit and there is no internal browning of the fruit. This disorder is more often seen during cool, wet and cloudy conditions. It is worsened by too much or too little water. High N and/or low K will cause an increase in the disorder. Older varieties are often more susceptible to this disorder.

Research in California indicates that for proper fruit color development higher K levels than are necessary for yield alone are needed. Soils and plants with high K had lower levels of the disorder. Foliar applications of K were not totally effective in reducing the disorder. Work in Michigan suggests that soils high in organic matter (above 3.5%) helped to reduce the disorder in a tomato crop. In addition, soils with a pH of 6.4 had low incidence of yellow

shoulder while tomatoes grown on soils with a pH above 6.7 had a high incidence.

Growers should have K tissue levels of at least 3% before fruit is one inch in diameter. In addition, the ratio of Magnesium (Mg) to Calcium (Ca) is important and a ratio of Mg:Ca of 1:4 to 1:6 should be maintained in the crop.

*Graywall* appears as grayish and sometimes sunken areas on a fruit. Internally the vascular tissue is brown resulting from collapse of the tissue. This can occur on the outer part of the fruit as well as in the center. It is usually more of a problem with cool, short days and often occurs in a late tomato crop. Graywall usually develops in green fruit but can occur as fruit is ripening. Fruit do not ripen properly and will have a blotchy appearance making them unmarketable. Graywall occurs on any part of the fruit. High N may increase the incidence of graywall and adequate K may reduce the problem. The disorder may also be caused by stress on the plants resulting from drought, excessive heat, root problems, severe nutrient deficiencies, etc. and there are varietal differences in susceptibility. This disorder is not clearly understood. Note that internal browning can also be caused by tobacco mosaic virus (TMV).

*Internal white tissue* is a disorder where the fruit usually show no external symptoms. When a ripe, affected fruit is cut there will be white, hard areas found in the outer tissue and sometimes in the center of the fruit as well. High temperatures during ripening are believed to be the cause of this disorder. Maintaining adequate K in the soil may reduce but not eliminate it. Some varieties are more susceptible to this disorder, especially high colored varieties. This disorder can be severe enough to cause fruit to be unmarketable.

### **Catfacing**

Catfacing is where fruit are malformed and scarred, usually at the blossom end. It is caused by exposure of seedlings to 60-65°F (16-18°C) day temperatures and 50-60°F (10-16°C) night temperatures for 1 week, approximately 4 weeks before pollination. The first flower cluster is susceptible to low temperature-induced catfacing when seedlings have 4-5 true leaves. Fruit on later clusters will show catfacing if exposed to low temperatures in the field. Avoid hardening seedlings by exposure to low temperatures. Varieties differ in their susceptibility to the disorder.

### **Cracking**

Cracking is due to the rapid uptake of water, resulting in enlargement of cells and separation of the epidermis of the fruit. Water can be taken up through the roots or through the tissue around the stem scar. The type of cracking (concentric, radiating out from the stem, or diagonal or transverse cracks across the fruit) is determined primarily by fruit structure and variety. Different types of cracking may be present in a variety or an individual fruit.

The severity of cracking is determined by water availability, variety and maturity. As the fruit ripens, the bonding between cells progressively weakens, resulting in more severe cracking. High rainfall and irrigation, or frequent low to moderate rainfall, especially following a period of low soil moisture may increase cracking. To minimize cracking, select a crack-resistant variety, maintain a high calcium level in the soil and keep fruit growing at a uniform rate by maintaining uniform soil moisture levels. Maintain good fruit cover by proper fertilization and fungicide applications. Harvest fruit at the earliest stage of maturity that is acceptable by your market.

### **Russetting**

Russetting, or weather checking of the surface of the fruit is caused by the presence of water on the fruit surface for extended periods of time when there are frequent light rainfalls, mist, fog, and dew. Wide fluctuations in temperature of exposed fruit also contribute to this disorder. Russetting can cause fruit to be unmarketable. Maintain good fruit cover by proper fertilization and fungicide applications. Use varieties that feature a dense canopy and resistance to foliar diseases.

### **Sunburn and Sunscald**

Sunburn and sunscald result from exposure to direct sunlight. Mild sunburn appears as yellowish or yellow-red color of fruit on the side exposed to the sun. Severe symptoms appear as whitish, water-soaked, scalded, or blistered areas. Sunscald is more severe on fruit that developed in shaded conditions but was exposed to direct sunlight after defoliation or harvesting. Under dry conditions, the white areas can become dry and leathery. Secondary infection can produce a dark, dry rot. Under moist conditions, scalded areas can decay from secondary infections. To control sunburn and sunscald, select varieties with good fruit cover, supply sufficient water and nutrients to provide good vegetative growth and manage pests. Train workers to avoid turning vines during harvesting or to reposition vines to shade fruit.

### **Yellow Shoulders**

Yellowing may occur on the shoulders of fruit exposed to the sun, especially on varieties that have darker green shoulders when immature (those lacking the uniform ripening gene). The tissue beneath the yellow shoulder is usually corky and may vary from greenish white to pale yellow. Select varieties with the uniform ripening gene and provide good fruit cover as described above.



## Weed Control

### THE LABEL IS THE LAW-see the Pesticide Use Disclaimer on the first page of chapter F. Recommended Herbicides

1. Identify the weeds in each field and select recommended herbicides. More information is available in the “Herbicide Effectiveness on Common Weeds in Vegetables” (Table E-2) in chapter E Pest Management.
2. Minimize herbicide resistance development. Identify the herbicide site of action group number and follow recommended good management practices; **bolded group numbers in tables below are herbicides at higher risk for selecting resistant weed populations.** Include non-chemical weed control whenever possible.

Labeled Applications Sites for Tomato									
Herbicides	WSSA group number	Plastic mulch production					Bare-ground production		
		Soil-Applied		Postemergence			Soil-applied	POST	Post-harvest
		Under Plastic	Row Middles	Over Plastic	Row Middles	Post-Harvest			
Sandea	2	YES***	YES		YES		YES**	YES	
Dacthal	3							YES#	
Prowl H2O	3		YES				YES***		
Treflan	3		YES				YES***		
Metribuzin	5	YES	YES		YES		YES	YES	
Reflex*	14	YES	YES		YES		YES***		
Devrinol	15	YES	YES				YES		
Dual	15	YES	YES				YES		
Select	1			YES				YES	
SelectMax	1			YES				YES	
Poast	1			YES				YES	
Matrix	2		YES		YES			YES	
Gramoxone*	22				YES	YES			YES
Reglone*	22				YES	YES			YES

\*Special Local Needs Label 24(c), be sure it is registered for the specific state and for the intended use.

\*\*Delay transplanting for 7 days after application; not labeled for direct-seeding. \*\*\*Transplants only.

#Dacthal is labeled for over the top application, but will it will not control emerged weeds.

### 1. Soil Applied

Group	Product Name	Product Rate	Active Ingredient (*=Restricted Use)	Active Ingredient Rate	PHI (d)	REI (h)
2	Sandea 75DF	0.5 to 1.0 oz/A	halosulfuron	0.023 to 0.047 lb/A	30	12
<p><b>-Plasticulture:</b> under plastic application is labeled delay transplanting 7 days after herbicide application. Apply in a band under the plastic, immediately before laying the mulch; use on transplants only (not for seeded tomatoes), avoid herbicide treated soil from moving into the holes during transplanting. Plasticulture: labeled for row middle application with directed/shield application.</p> <p><b>-Bareground:</b> for transplants only: apply preplant incorporated 7 days before transplanting; use on transplants only (not for seeded tomatoes), avoid herbicide treated soil from moving into the holes during transplanting.</p> <p>-Bareground: for directed-seeded apply as directed/shielded application to row middles</p> <p>-Suppresses or controls yellow nutsedge and certain broadleaf weeds. Sandea provides both residual and postemergence control of susceptible weed species. Effective postemergence control requires an adjuvant.</p> <p>-Sandea is an ALS inhibiting herbicide and resistant weed populations are common in the region. <b>Do not</b> use Group 2 herbicides repeatedly in the same field. <b>Do not</b> apply Sandea to crops treated with a soil applied organophosphate insecticide, or use a foliar applied organophosphate insecticide within 21 days before or 7 days after a Sandea application.</p> <p>-Maximum Sandea applications per year is 2 and <b>do not</b> exceed 2 oz/A during the crop season.</p>						
3	Dacthal 6F Dacthal W-75	8.0 to 14.0 pt/A 6.0 to 14 lb/A	DCPA	6.0 to 10.5 lb/A	--	12
<p>-Labeled for applications over the top of transplants without injury (will not control emerged weeds); transplants should be well-established and growing conditions favorable for good plant growth.</p> <p>-Label recommends 4 to 6 weeks after transplanting or direct-seeded plants at 4 to 6 inches in height</p> <p>-Post-transplant applications can only be made with bare-ground production.</p> <p>-Dacthal will not control emerged weeds; apply to weed-free soils.</p> <p>-Primarily controls annual grasses and a few broadleaf weeds, including common purslane.</p> <p>-Results have been most consistent when used in fields with coarse -textured soils low in organic matter, and when the application are followed by rainfall or irrigation. -Maximum application not addressed on label.</p>						

1. Soil Applied - continued on next page

## 1. Soil Applied - continued

3	Prowl H2O 3.8CS	1.0 to 3.0 pt/A	<b>pendimethalin</b>	0.48 to 1.42 lb/A	70	24
<p><b>-Plasticulture:</b> recommended for row middles only. Labeled for under plastic, but no local data or experience with this application.  <b>-Bareground:</b> broadcast preplant or preplant incorporated before transplanting; not labeled for direct-seeded crop.  -Avoid root contact with Prowl-treated soil when placing transplants into furrow or hole or injury may occur.  -Prowl labeled for directed application to transplanted or established direct-seeded peppers; avoid contact with leaves or stems.  -Prowl will not control emerged weeds, only provides residual control; row middle applications may be made with Gramoxone using shielded sprayers. Use the lower rate on coarse-textured or sandy soils. Activate with ½ inch of rainfall or sprinkler irrigation within 48 hr of application to control most annual grasses and certain broadleaf weeds. -Maximum Prowl H2O application per season is 3 pt/A.</p>						
3	Treflan 4E	1 to 2 pt/A	<b>trifluralin</b>	0.5 to 1.0 lb/A	--	12
<p><b>-Plasticulture:</b> labeled for row middles only. <b>-Bareground:</b> broadcast preplant or preplant incorporated before transplanting; not labeled for direct-seeded crop. All applications need to be mechanically incorporating.  -Stunting may occur if weather is cool and damp at time of transplanting. Maximum application per season: not specified.</p>						
5	Metribuzin 75DF Metribuzin 4L	0.33 to 0.66 lb/A 0.5 to 1 pt/A	<b>metribuzin</b>	0.25 to 0.5 lb/A	7	12
<p><b>-Plasticulture:</b> under plastic application is labeled; apply in a band under the plastic, immediately before laying the mulch; use on transplants only (not for seeded tomatoes), roots of the transplants need to be placed below the zone of treated soil. There is no local data and limited experience with this use. Plasticulture: labeled for row middle application with directed/shield application.  <b>-Bareground:</b> broadcast preplant or preplant incorporated before transplanting; use on transplants only (not for seeded tomatoes), roots of the transplants need to be placed below the zone of treated soil.  -Metribuzin primarily controls broadleaf weeds and is weak on grasses; tankmix to improve grass control.  -Metribuzin has some postemergence activity. To get consistent control, apply metribuzin before weeds are 1 inch tall.  -Rainfastness is 6 h. -Maximum for metribuzin 75DF: 1.33 lb/A per crop season; metribuzin 4L: 2 pt/A per crop season.</p>						
14	Reflex 2SL	16 to 24 fl oz/A VA	<b>fomesafen</b>	0.25 to 0.375 lb/A	70	24
<p><b>-Special Local Needs Label 24(c) has been approved for VA (expires 12/31/2020).</b> -The use of Reflex 2SL is legal ONLY if a waiver of liability has been completed (see <a href="http://www.syngenta-us.com/labels/indemnified-label-login">www.syngenta-us.com/labels/indemnified-label-login</a>).  <b>-Only labeled for transplanted tomatoes; do not</b> use on directed seeded crop. Ensure that Reflex treated soil is not moved into the transplant holes.  <b>-Plasticulture:</b> under plastic application is labeled; apply in a band under the plastic, immediately before laying the mulch; <b>do not</b> mechanically incorporate. Crops may be transplanted immediately following application.  -Plasticulture: labeled for application over the top of plastic before transplanting only if beds are shaped to allow herbicide to be readily washed off with irrigation or rainfall; and single rainfall or irrigation provides at least 0.5 inches of water before transplanting; and plastic does not have any holes until after Reflex has been washed off.  -Plasticulture: labeled row middles application prior to transplanting.  <b>-Bareground:</b> labeled for pre-transplant applied to soil surface, <b>do not</b> mechanically incorporate. Rainfall or irrigation between herbicide application and transplanting will likely reduce the risk of crop injury due to splashing.  -Reflex provides both residual and postemergence control of susceptible weed species. Effective postemergence control requires an adjuvant. Varieties may vary in their response to Reflex; treat small acreages first to determine crop tolerance.  -Consider rotational crops when applying fomesafen. If crop is replanted <b>do not</b> re-apply. Rotational restrictions depend on whether fomesafen was applied bareground, or under or over plastic mulch, see 24(c) label for specifics.  -Maximum Reflex application: VA 24 fl oz/A IN ALTERNATE YEARS.</p>						
15	Devrinol 2-XT 2EC Devrinol DF-XT 50DF	2 to 4 qt/A 2 to 4 lb/A	<b>napropamide</b>	1.0 to 2.0 lb/A	--	24
<p><b>-Plasticulture:</b> under plastic is labeled for seeded or transplanted tomatoes; apply in a band under the plastic, immediately before laying mulch. Use lower rate on coarse textured or sandy soil. Condensation that forms on the underside of the mulch will activate the herbicide. Plasticulture: row middles application is labeled.  <b>-Bareground:</b> apply as broadcast, preemergence treatment for seeded and transplanted tomatoes. Rainfall or irrigation within 24 hr after application improves performance (½ inch sprinkler irrigation).  -Annual grasses and certain annual broadleaf weeds will be suppressed or controlled. May reduce stand and yield of fall planted small grain crop. Moldboard plowing will reduce the risk of injury.  -Maximum Devrinol application per season: 4 qt/A (2-XT) or 4 lb/A (DF-XT).</p>						
15	Dual Magnum 7.62E	1.0 to 2.0 pt/A	<b>s-metolachlor</b>	0.95 to 1.9 lb/A	30 to 90	24
<p><b>-Plasticulture:</b> under plastic is labeled transplanted tomatoes; apply in a band under the plastic, immediately before laying mulch. Use lower rate on coarse textured or sandy soil. Condensation that forms on the underside of the mulch will activate the herbicide. Plasticulture: row middles application is labeled.  <b>-Bareground:</b> apply for preplant incorporated or broadcast, preemergence treatment before transplanting tomatoes. Seeded tomatoes can be treated when at least 4 inches tall at time of application and spray is directed at the soil and minimal amounts of herbicide contact tomato plants. Avoid moving treated soil into transplant holes.  -Use lower rates on coarse-textured soils low in organic matter and higher rates on fine-textured soils with greater organic matter.  -Application to varieties with unknown tolerance to Dual Magnum may result in crop injury. Transplants weakened by any cause may be injured by Dual Magnum. Plant healthy transplants and avoid planting when wet, cool, or unfavorable growing conditions exist.  -Delaying transplanting for 7 days or more can reduce risk of injury.  <b>-Do not harvest</b> within 90 days of application if more than 1.33 pt/A was used per season; PHI is 30 days if 1.33 pt/A or less is used. <b>For VA only</b> PHI is 60 days if 1.67 pt/A per season or less was applied; PHI is 90 days if more than 1.67 pt/A per season was used (VA 24(c) expires 12/31/2020). <b>Do not</b> exceed 2 applications per growing season.</p>						

F Tomatoes

2. Postemergence						
Group	Product Name	Product Rate	Active Ingredient (*=Restricted Use)	Active Ingredient Rate	PHI (d)	REI (h)
1	Select 2EC	6 to 8 fl oz/A	clethodim	0.07 to 0.12 lb/A	20	24
	Select Max 0.97EC	9 to 16 fl oz/A				
	Poast 1.5EC	1 to 2.5 pt/A	sethoxydim	0.2 to 0.5 lb/A	20	12
<p><b>-Select 2EC:</b> use crop oil concentrate (COC) at 1% v/v (1 gal/100 gal of spray solution). <b>Select Max:</b> use nonionic surfactant (NIS) at 0.25% v/v (1 qt/100 gal of spray solution). <b>Poast:</b> Apply with COC at 1.0% v/v</p> <p><b>-The use of COC may increase the risk of crop injury when hot or humid conditions prevail.</b> To reduce the risk of crop injury, omit additives or switch to NIS when grasses are small and soil moisture is adequate.</p> <p>-Use lower labeled rates for annual grass control and higher labeled rates for perennial grass control.</p> <p>-Yellow nutsedge, wild onion, wild garlic, and broadleaf weeds will not be controlled.</p> <p>-Controls many annual and certain perennial grasses, including annual bluegrass, but Poast is preferred for goosegrass control. For best results, treat annual grasses when they are actively growing and before tillers are present. Control may be reduced if grasses are large or under hot or dry weather conditions. Repeated applications may be necessary to control certain perennial weeds. If repeat applications are necessary, allow 14 days between applications. Rainfastness is 1 h.</p> <p><b>-Do not</b> tank-mix with or apply within 2 to 3 days of any other pesticide, unless labeled, as this may increase the risk of crop injury or reduce the control of grasses. <b>-Do not</b> apply more than 8 fl oz of Select 2EC in a single application and <b>do not</b> exceed 2 pt/A for the season; <b>do not</b> apply more than 16 fl oz of Select Max in a single application and <b>do not</b> exceed 4 pt/A for the season.</p> <p><b>-Do not</b> apply more than 1.5 pt/A Poast 1.5EC in single application and <b>do not</b> exceed 4.5 pt/A for the season.</p>						
2	Matrix 25DF	1.0 to 2.0 oz/A	rimsulfuron	0.0156 to 0.0312 lb/A	45	4
	Solida 25DF					
<p>-Apply early postemergence but not before the crop has at least 2 full-sized true leaves (label allows applications as early as cotyledon stage of tomatoes; but no local data is available at that stage). Not recommended for over the top application with plasticulture.</p> <p>-Apply with nonionic surfactant at 0.25% v/v (1.0 qt/100 gal of spray solution); use of an adjuvant may cause temporary chlorosis, but symptoms usually disappear within 5 to 15 days.</p> <p>-Controls many weeds including foxtail species, pigweed species, wild mustard, and wild radish. Suppresses common lambsquarters, common ragweed, jimsonweed, morningglory species, and yellow nutsedge. Optimum performance is obtained when weeds are less than 1 inch in height and are actively growing. Tank mix with metribuzin to improve broadleaf weed control.</p> <p>-Best results occur with 0.5 inches of rainfall or irrigation no sooner than 4 h but not more than 5 days after application.</p> <p>-Matrix provides both residual and postemergence control of susceptible weed species.</p> <p>-Matrix is an ALS inhibiting herbicide and resistant weed populations are common in the region. <b>Do not</b> use Group 2 herbicides repeatedly in the same field. -Rainfastness is 4 h. -Maximum for Matrix: 4 oz/A per year.</p>						
2	Sandea 75DF	0.5 to 1.0 oz/A	halosulfuron	0.023 to 0.047 lb/A	30	12
<p>-Apply over the top, post directed, or with crop shields; not recommended for over the top application with plasticulture.</p> <p>-Apply to tomato plants that are established, actively growing and a minimum of 14 days after transplanting or after the 4<sup>th</sup> leaf stage of seeded tomatoes. Applications during bloom can cause bloom drop under certain environmental conditions.</p> <p>-Apply with nonionic surfactant at 0.25% v/v (1.0 qt/100 gal).</p> <p>-Provides control of yellow nutsedge and certain annual broadleaf weeds. Control of weeds taller than 3 inches may not be adequate.</p> <p>-Sandea provides both residual and postemergence control of susceptible weed species.</p> <p>-Sandea is an ALS inhibiting herbicide and resistant weed populations are common in the region. <b>Do not</b> use Group 2 herbicides repeatedly in the same field. <b>Do not</b> apply Sandea to crops treated with a soil-applied organophosphate insecticide, or use a foliar applied organophosphate insecticide within 21 days before or 7 days after a Sandea application. -Rainfastness is 4 h.</p> <p><b>-Do not</b> apply more than 2 applications, or more than 2 oz of product, per crop cycle; <b>do not</b> exceed 2 oz/A per 12 month period.</p>						
3	Dacthal 6F	8.0 to 14.0 pt/A	DCPA	6.0 to 10.5 lb/A	--	12
	Dacthal W-75	6.0 to 14 lb/A				
<p>-Labeled for applications over the top of transplants.</p> <p>-Dacthal will not control emerged weeds; apply to weed-free soils. See comments under soil applied section</p>						
5	Metribuzin 75DF	0.33 to 0.66 lb/A	metribuzin	0.25 to 0.5 lb/A	7	12
	Metribuzin 4L	0.5 to 1 pt/A				
<p>-Apply over the top, post directed, or with crop shields; not recommended for over the top application with plasticulture. Maximum rate for over the top application is 0.67 lb (75DF) or 1 pt (4L); and maximum rate for post directed is 1.33 lb (75DF) or 2 pt (4L).</p> <p>-Apply postemergence to transplants with at least 5 true leaves and have recovered from transplant shock (new growth evident) or at least 2 weeks after transplanting. Transplant with fewer than 5 true leaves are at greater risk of herbicide injury.</p> <p><b>-Do not</b> use hot caps on tomatoes within 7 days before or after application. <b>-Do not</b> apply within 3 days after periods of cool, wet, or cloudy weather or crop injury will occur. <b>-Do not</b> apply within 24 h of applications of other pesticides.</p> <p>-Allow at least 14 days between applications or severe crop injury may occur.</p> <p>-Metribuzin primarily controls broadleaf weeds and is weak on grasses.</p> <p>-Metribuzin has some postemergence activity. To get consistent control, apply metribuzin before weeds are 1 inch tall.</p> <p>-Tank mix with appropriate postemergence herbicides if weeds are emerged at time of application.</p> <p>-Rates up to 1.3 lb of metribuzin 75DF or 2 pt of metribuzin 4L are labeled for directed applications, refer to label.</p> <p>-Maximum for metribuzin 75DF: 1.33 lb/A per crop season; metribuzin 4L: 2 pt/A per crop season.</p>						

2. Postemergence - continued on next page

## 2. Postemergence - continued

22	Reglone 2SL	1 qt/A	<b>diquat</b>	0.5	30	24
-A Special Local Needs Label 24(c) has been approved in NJ (expires 12/31/2021). -Apply as post-directed application to the row middles either prior to transplanting or with a hooded sprayer to row middles when transplants are well established. Do not allow spray to contact crop foliage. -Always include non-ionic surfactant at 2 pt/100 gal. Spray coverage is essential for optimum effectiveness. -Rainfastness 30 min. A maximum of 2 applications during the growing season are allowed.						
22	Gramoxone 2SL	2.0 pt/A	<b>paraquat*</b>	0.5 lb/A	30	24
-Gramoxone can be applied before or after transplanting to control emerged broadleaf weeds and grass seedlings. -Include a nonionic surfactant at 0.25% v/v. <b>Do not</b> allow spray to contact crop foliage as injury may result. Use flaps that drag along the edge of plastic mulch and use low spray pressure (maximum of 30 psi) to reduce small droplets that are prone to drift. -See the label for additional information and warnings. -Rainfastness is 30 min. A maximum of 3 applications per year are allowed. - <b>Restricted-use pesticide.</b> Only certified applicators, who successfully complete the paraquat-specific training, can mix, load or apply paraquat. Application of paraquat "under the direct supervision" of a certified applicator is no longer allowed. Required training link ( <a href="http://usparaquattraining.com">http://usparaquattraining.com</a> ); certified applicators must repeat training every three years.						

## 3. Postharvest

Group	Product Name	Product Rate	Active Ingredient (*=Restricted Use)	Active Ingredient Rate	PHI (d)	REI (h)
22	Reglone 2SL	1.5 pt/A	<b>diquat</b>	0.375	--	24
-A Special Local Needs Label 24(c) has been approved in NJ (expires 12/31/2021) for postharvest application to desiccate the crop. -Apply after the last harvest for bareground or plasticulture. Always include non-ionic surfactant at 1 to 2 pt/100 gal. -Spray coverage is essential for optimum effectiveness, label recommends 60 to 100 gal/A. -Rainfastness 30 min.						
22	Gramoxone 2SL	2.4 to 3.75 pt/A	<b>paraquat*</b>	0.6 to 0.94 lb/A	--	24
-Apply after the last harvest for bareground or plasticulture. Always include an adjuvant. -Spray coverage is essential for optimum effectiveness. See the label for additional information and warnings. -Rainfastness 30 min. A maximum of 2 applications for crop desiccation are allowed. - <b>Restricted-use pesticide.</b> Only certified applicators, who successfully complete the paraquat-specific training, can mix, load or apply paraquat. Application of paraquat "under the direct supervision" of a certified applicator is no longer allowed. Required training link ( <a href="http://usparaquattraining.com">http://usparaquattraining.com</a> ); certified applicators must repeat training every three years.						

## 4. Other Labeled Herbicides These products are labeled but limited local data are available; and/or are labeled but not recommended in our region due to potential crop injury concerns.

Group	Product Name	Active Ingredient (*=Restricted Use)
2	Envoke	<b>trifloxysulfuron</b>
2	League	<b>imazosulfuron</b>
14	Aim	<b>carfentrazone</b>
14	Vida	<b>pyraflufen</b>
14	Spartan	<b>sulfentrazone</b>

## Insect Control

**THE LABEL IS THE LAW-see the Pesticide Use Disclaimer on the first page of chapter F. Recommended Insecticides Field Tomatoes (Fresh Market and Processing Tomatoes)**

## Aphids

Tomatoes in the mid-Atlantic U.S. can suffer late-season infestations from green peach aphids or potato aphids. Frequent pyrethroid applications are often the cause of outbreaks of green peach aphid. Adequate coverage of the undersides of leaves is important for effective aphid control.

Apply one of the following formulations (thorough spray coverage between leaves is important):						
Group	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
1A	Lannate LV	1.5 to 3.0 pt/A	methomyl*	3	48	H
4A	Neonicotinoid insecticides registered for use on Tomatoes: see table at the end of Insect Control.					
4C	Closer SC	1.5 to 2.0 fl oz/A	sulfoxaflor	1	12	H
4C	Transform WG	0.75 to 1.0 oz/A	sulfoxaflor	1	24	H
4D	Sivanto Prime or 200SL	21.0 to 28.0 fl oz/A	flupyradifurone - <b>soil</b>	45	4	M
4D	Sivanto Prime or 200SL	10.5 to 14.0 fl oz/A	flupyradifurone - <b>foliar</b>	1	4	M
9B	Fulfill 50WDG	2.75 oz/A	pymetrozine	0	12	L

Aphids - continued on next page

## F Tomatoes

### Aphids - continued

9B	PQZ	2.4 to 3.2 fl oz/A	pyrifluquinazon	1	12	L
9D	Sefina	3.0 fl oz/A	afidopyropen	0	12	L
21A	Torac	17.0 to 21.0 fl oz/A	tolfenpyrad	1	12	H
23	Movento	4.0 to 5.0 fl oz/A	spirotetramat	1	24	L
28 + 6	Minecto Pro	10.0 fl oz/A	cyantraniliprole + abamectin*	7	12	H
29	Beleaf 50SG	2.8 to 4.3 oz/A	flonicamid	0	12	L

### Caterpillar “Worm” Pests Including: Corn Earworms (=Tomato Fruitworms) (CEW), European Corn Borers (ECB), Beet Armyworms (BAW), Cabbage Loopers (CL), Hornworms, and other Armyworms

Tomatoes are frequently attacked by various lepidopteran pest species. CEW (or tomato fruitworm) is the most important pest species, but a mix of any of the above-listed pest species can also contribute to “worm” damage on tomatoes. Local pheromone or blacklight traps are effective for monitoring key moth pest populations. Consult your Extension Agent or IPM alerts for information about trap catches. Also, visually inspecting plants and fruit or beat sheeting can help determine the presence or absence of lepidopteran pests. There is no reliable economic threshold. Note that not all lepidopteran pest species are listed on all of the insecticide labels below, but, unless noted, these products have activity on all caterpillars. **Pyrethroid (Group 3A) resistance is common in BAW and also has been reported in populations of CEW. So caution should be used when using that class of insecticide. Rotating insecticide classes within a season is strongly recommended.**

Apply one of the following formulations:						
Group	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
1A	Lannate LV	1.5 to 3.0 pt/A	methomyl*	3	48	H
3A	Pyrethroid insecticides registered for use on Tomatoes: see table at the end of Insect Control.					
4A	Neonicotinoid insecticides registered for use on Tomatoes: see table at the end of Insect Control.					
5	Entrust SC (OMRI)	3.0 to 6.0 fl oz/A	spinosad	1	4	M
5	Radiant SC	5.0 to 10.0 fl oz/A	spinetoram	1	4	M
6	Proclaim 5SG	2.4 to 4.8 oz/A	emamectin benzoate*	7	12	H
11A	Dipel DF, others (OMRI)	1.0 to 2.0 lb/A	<i>Bacillus thuringiensis kurstaki</i>	0	4	N
11A	XenTari (OMRI)	0.5 to 2.0 lb/A	<i>Bacillus thuringiensis aizawai</i>	0	4	N
15	Rimon 0.83EC	9.0 to 12.0 fl oz/A	novaluron	1	12	M
18	Confirm 2F	6.0 to 8.0 fl oz/A (early season); 8.0 to 16.0 fl oz/A (late season)	tebufenozide (not labeled for CEW)	7	4	M
18	Intrepid 2F	4.0 to 8.0 fl oz/A (early season); 8.0 to 16.0 fl oz/A (late season) (ECB, HW, CL only)	methoxyfenozide	1	4	L
22	Avaunt 30WDG, Avaunt eVo	2.5 to 3.5 oz/A (HW, CL); 3.5 oz/A (CEW)	indoxacarb	3	12	H
28	Coragen 1.67SC	3.5 to 7.5 fl oz/A	chlorantraniliprole - soil	1	4	L
28	Coragen 1.67SC	3.5 to 7.5 fl oz/A	chlorantraniliprole - foliar	1	4	L
28	Exirel	7.0 to 13.5 fl oz/A (CEW, HW, ECB); 10.0 to 17.0 fl oz/A (CL)	cyantraniliprole	1	12	H
28	Verimark	5.0 to 10.0 fl oz/A (CEW, HW); 6.75 to 10.0 fl oz/A (CL)	cyantraniliprole	1	4	H
28 + 6	Minecto Pro	7.5 to 10.0 fl oz/A	cyantraniliprole + abamectin*	7	12	H

### Colorado Potato Beetles (CPB)

Rotation to crops other than potato, tomato, and eggplant is extremely important in reducing CPB problems. Also, transplants placed into no-till fields, mulches or other crop residue will reduce or delay potato beetle infestations. Look for CPB adults shortly after seedling emergence or transplanting. Early season populations tend to be concentrated in areas where tomatoes or potatoes were previously grown. Thoroughly scout tomato fields and spray only when necessary. Also spot treatment of "hot spots" only is recommended if infestation is localized. For established direct-seeded or transplant tomatoes, begin treatment if the population level exceeds 15 CPB adults per 10 plants throughout the field. If early treatment is not applied, wait for egg hatch and spray when larvae are young and exceed 20 CPB larvae and/or adults per 10 plants. Reassess after each treatment. Avoid the application of late-season sprays to prevent the buildup of insecticide-resistant beetles.

*Colorado Potato Beetles (CPB) - continued on next page*

## Colorado Potato Beetles (CPB) - continued

Apply one of the following formulations:						
Group	Product Name	Product Rate	Active Ingredient(s) (* = Restricted Use)	PHI (d)	REI (h)	Bee TR
1A	Vydate L	2.0 to 4.0 pt/A	oxamyl* - foliar	7	48	H
4A	Neonicotinoid insecticides registered for use on Tomatoes: see table at the end of Insect Control.					
4D	Sivanto Prime or 200SL	10.5 to 14.0 fl oz/A	flupyradifurone - foliar	1	4	M
5	Entrust SC (OMRI)	3.0 to 6.0 fl oz/A	spinosad	1	4	M
5	Radiant SC	5.0 to 10.0 fl oz/A	spinetoram	1	4	M
6	Agri-Mek SC	1.75 to 3.5 fl oz/A	abamectin*	7	12	H
6 + 3A	Gladiator	19.0 fl oz/A	abamectin* + zeta-cypermethrin*	7	12	H
11A	Trident (OMRI)	3.0 to 6.0 qt/A	<i>Bacillus thuringiensis tenebrionis</i>	0	4	L
15	Rimon 0.83EC	9.0 to 12.0 fl oz/A	novaluron	1	12	M
28	Coragen 1.67SC	3.5 to 7.5 fl oz/A	chlorantraniliprole - soil	1	4	L
28	Coragen 1.67SC	3.5 to 7.5 fl oz/A	chlorantraniliprole - foliar	1	4	L
28	Exirel	7.0 to 13.5 fl oz/A	cyantraniliprole	1	12	H
28	Verimark	5.0 to 10.0 fl oz/A	cyantraniliprole	1	4	H
28	Harvanta 50SL	10.9 to 16.4 fl oz/A	cyclaniliprole	1	4	H
28 + 6	Minecto Pro	5.5 to 10.0 fl oz/A	cyantraniliprole + abamectin*	7	12	H

**Cutworms** See also section E 3.1. Soil Pests - Detection and Control.

Apply one of the following formulations:						
Group	Product Name	Product Rate	Active Ingredient(s) (* = Restricted Use)	PHI (d)	REI (h)	Bee TR
<b>Preplanting Field Treatment. Just before seeding or transplanting, broadcast on the soil surface the following:</b>						
1B	Diazinon AG500	2.0 to 4.0 qt/A	diazinon*	n/a	48	H
3A	Capture LFR	3.4 to 6.8 fl oz/A	bifenthrin*	n/a	12	H
<b>Postplanting Treatment. If control is required after seedling emergence or after transplanting, treat soil thoroughly beneath plants with the following:</b>						
3A	Pyrethroid insecticides registered for use on Tomatoes: see table at the end of Insect Control.					

**Flea Beetles**

Are small dark beetles that feed by chewing round holes into leaves. They usually only pose a problem to small tomato plants. Heavy feeding on small plants during hot dry periods can result in stand loss. Watch for flea beetle feeding on transplanted tomatoes. If needed one insecticide application is usually all that is needed for control.

Apply one of the following formulations:						
Group	Product Name	Product Rate	Active Ingredient(s) (* = Restricted Use)	PHI (d)	REI (h)	Bee TR
3A	Pyrethroid insecticides registered for use on Tomatoes: see table at the end of Insect Control.					
4A	Neonicotinoid insecticides registered for use on Tomatoes: see table at the end of Insect Control.					
21A	Torac	17.0 to 21.0 fl oz/A	tolfenpyrad	1	12	H
28	Verimark	6.75 to 13.5 fl oz/A	cyantraniliprole	1	4	H
28	Harvanta 50SL	10.9 to 16.4 fl oz/A	cyclaniliprole	1	4	H

**Leafminers**

Leafminers are generally not a significant problem in most fields. Adults are small, black and yellow flies that insert their eggs into leaves and the larvae feed between leaf surfaces, creating a meandering track. A few of these tracks per leaf will not affect yield.

Group	Product Name	Product Rate	Active Ingredient(s) (* = Restricted Use)	PHI (d)	REI (h)	Bee TR
<b>Treat with one of the following formulations when first mines appear and repeat every 7 days or as needed.</b>						
1B	Dimethoate 400	0.5 to 1.0 pt/A	dimethoate*	7	48	H
3A	Pyrethroid insecticides registered for use on Tomatoes: see table at the end of Insect Control.					
4A	Neonicotinoid insecticides registered for use on Tomatoes: see table at the end of Insect Control.					
5	Entrust SC (OMRI)	6.0 to 10.0 fl oz/A	spinosad	1	4	M
5	Radiant SC	6.0 to 10.0 fl oz/A	spinetoram	1	4	M
6	Agri-Mek SC	1.75 to 3.5 fl oz/A	abamectin*	7	12	H
15	Rimon 0.83EC	12 fl oz/A	novaluron	1	12	M
17	Trigard 75WSP	2.66 oz/A	cyromazine	0	12	H

Leafminers - continued on next page

## F Tomatoes

### Leafminers - continued

28	Coragen 1.67SC	5.0 to 7.5 fl oz/A	chlorantraniliprole - <b>soil</b>	1	4	L
28	Coragen 1.67SC (larvae only)	5.0 to 7.5 fl oz/A	chlorantraniliprole - <b>foliar</b>	1	4	L
28	Exirel	13.5 to 20.5 fl oz/A	cyantraniliprole	1	12	H
28	Verimark	6.75 to 13.5 fl oz/A	cyantraniliprole ( <b>at planting</b> )	1	4	H
28	Harvanta 50SL	10.9 to 16.4 fl oz/A	cyflumetofen	1	4	H
28 + 6	Minecto Pro	5.5 to 10.0 fl oz/A	cyantraniliprole + abamectin*	7	12	H

## Mites

Mite infestations generally begin around field margins, grassy areas, and windbreaks during hot dry periods that favor the mites. Beginning infestations sometime occur from infested transplants that were produced in GHs with bedding plants-be sure to check transplants for mites. **DO NOT** mow or maintain grassy areas after mid-summer since this forces mites into the crop. Localized infestations can be spot treated. Watch for mite feeding, *i.e.*, stippling of leaves, in mid-summer. The use of dimethoate for other pests can reduce spider mite populations.

Apply one of the following formulations: Note: Thorough spray coverage beneath leaves is important.						
Group	Product Name	Product Rate	Active Ingredient(s) (*= <b>Restricted Use</b> )	PHI (d)	REI (h)	Bee TR
6	Agri-Mek SC	1.75 to 3.5 fl oz/A	abamectin*	7	12	H
6 + 3A	Gladiator	19.0 fl oz/A	abamectin* + zeta-cypermethrin*	7	12	H
10A	Onager 1EC	12 to 24 fl oz/A	hexythiazox	1	12	N
20B	Kanemite 15SC	31 fl oz/A	acequinocyl	1	12	L
21A	Portal XLO	2.0 pt/A	fenpyroximate	1	12	L
21A	Torac (broadmites only)	14.0 to 21.0 fl oz/A	tolfenpyrad	1	12	H
21A	Magister SC	24.0 to 36.0 fl oz/A	fenazaquin	3	12	H
23	Oberon 2SC	7.0 to 8.5 fl oz/A	spiromesifen	1	12	M
20D	Acramite 50WS	0.75 to 1.0 lb/A	bifenazate	3	12	M
25	Nealta	13.7 fl oz/A	cyflumetofen	3	12	--
28 + 6	Minecto Pro	5.5 to 10.0 fl oz/A	cyantraniliprole + abamectin*	7	12	H

## Pinworms

This pest is introduced on southern transplants. Begin sprays if leaf damage is observed. Late evening sprays may be most effective when moths are active.

Apply one of the following formulations:						
Group	Product Name	Product Rate	Active Ingredient(s) (*= <b>Restricted Use</b> )	PHI (d)	REI (h)	Bee TR
1A	Lannate LV	1.5 to 3.0 pt/A	methomyl*	1	48	H
3A	Pyrethroid insecticides registered for use on Tomatoes: see table at the end of Insect Control.					
4A	Neonicotinoid insecticides registered for use on Tomatoes: see table at the end of Insect Control.					
5	Entrust SC (OMRI)	4.0 to 8.0 fl oz/A	spinosad	1	4	M
5	Radiant SC	5.0 to 10.0 fl oz/A	spinetoram	1	4	M
6	Agri-Mek SC	3.5 fl oz/A	abamectin*	7	12	H
6	Proclaim 5SG	2.4 to 4.8 oz/A	emamectin benzoate*	7	12	H
6 + 3A	Gladiator	19.0 fl oz/A	abamectin* + zeta-cypermethrin*	7	12	H
15	Rimon 0.83EC	9.0 to 12.0 fl oz/A	novaluron	1	12	M
22	Avaunt 30WDG, Avaunt eVo	3.5 oz/A	indoxacarb	3	12	H
28	Coragen 1.67SC	3.5 to 7.5 fl oz/A	chlorantraniliprole - <b>soil</b>	1	4	L
28	Coragen 1.67SC (larvae)	3.5 to 7.5 fl oz/A	chlorantraniliprole - <b>foliar</b>	1	4	L
28	Exirel	7.0 to 13.5 fl oz/A	cyantraniliprole	1	12	H
28	Verimark	5.0 to 10.0 fl oz/A	cyantraniliprole	1	4	H
28	Harvanta 50SL	10.9 to 16.4 fl oz/A	cyflumetofen	1	4	H
28 + 6	Minecto Pro	10.0 fl oz/A	cyantraniliprole + abamectin*	7	12	H
n/a	NoMate TPW <sup>1</sup>	200 to 400 spirals/A	mating disruption hormone	n/a	n/a	n/a

<sup>1</sup>NoMate uses a disruption pheromone for preventing mating of emerging adults from young transplants. The pheromone is applied to a hard plastic matrix formed into a hanging "spiral" for dispersal into the air. Apply at first sign of pinworm larvae in leaves.

## Stink Bugs

Several different species of stink bugs feed on tomatoes, but they produce similar damage. Adult stink bugs are shield shaped and usually brown or green sometimes having colored markings. Adults overwinter on the ground

under leaves, or other protected areas. Feeding damage appears as dark pinpricks, surrounded by a white area that turns yellow on ripe fruit (cloudy spot). Because stinkbugs are so mobile and are quick to drop to the ground when the plant is disturbed there is no good scouting program for them. Watch the edges of fields for the first sign of cloudy spot to appear on green fruit. High spray gallonages and pressures are needed to penetrate the plant canopy to reach stinkbugs (especially immatures) that are hiding in the interior of the plant.

Apply one of the following formulations:						
Group	Product Name	Product Rate	Active Ingredient(s) (* = Restricted Use)	PHI (d)	REI (h)	Bee TR
1A	Lannate LV (brown mar-morated stink bug only)	3.0 pt/A	methomyl*	1	48	H
3A	Pyrethroid insecticides registered for use on Tomatoes: see table at the end of Insect Control.					
4A	Neonicotinoid insecticides registered for use on Tomatoes: see table at the end of Insect Control.					

## Thrips

Very high numbers of thrips can cause damage with their feeding, which distorts plant growth, deforms flowers, and causes small white marks (stippling) on emerging leaves that often have tiny black fecal specks in them. Several species of thrips also spread Tomato Spotted Wilt Virus. The virus can only be acquired by the immature stage of thrips while transmission primarily occurs via adults, which then transmit the virus their entire lives. Watch for the first signs of thrips which are stippling on leaves or thrips in flowers. Stippling marks or 3-5 thrips/flower indicate treatment may be necessary. Do not produce vegetable transplants with bedding plants in the same greenhouse.

Apply one of the following formulations:						
Group	Product Name	Product Rate	Active Ingredient(s) (* = Restricted Use)	PHI (d)	REI (h)	Bee TR
3A <sup>1</sup>	Pyrethroid insecticides registered for use on Tomatoes: see table at the end of Insect Control.					
4A	Neonicotinoid insecticides registered for use on Tomatoes: see table at the end of Insect Control.					
5	Entrust SC (OMRI)	4.0 to 8.0 fl oz/A	spinosad	1	4	M
5	Radiant SC	6.0 to 10.0 fl oz/A	spinetoram	1	4	M
21A	Torac	21.0 fl oz/A	tolfenpyrad	1	12	H
28	Harvanta 50SL	10.9 to 16.4 fl oz/A	cyclaniliprole	1	4	H

<sup>1</sup>Resistance concerns with Western flower thrips

## Whiteflies

Usually are only a problem late in the season on field tomatoes.

Apply one of the following formulations:						
Group	Product Name	Product Rate	Active Ingredient(s) (* = Restricted Use)	PHI (d)	REI (h)	Bee TR
4A	Neonicotinoid insecticides registered for use on Tomatoes: see table at the end of Insect Control.					
4C	Closer SC	4.25 to 4.5 fl oz/A	sulfoxaflor	1	12	H
4C	Transform WG	2.0 to 2.25 oz/A	sulfoxaflor	1	24	H
4D	Sivanto Prime or 200SL	21.0 to 28.0 fl oz/A	flupyradifurone - <b>soil</b>	45	4	M
4D	Sivanto Prime or 200SL	10.5 to 14.0 fl oz/A	flupyradifurone - <b>foliar</b>	1	4	M
7C	Knack	8.0 to 10.0 fl oz/A	pyriproxyfen	1	12	L
9B	Fulfill 50WDG	2.75 oz/A	pymetrozine	0	12	L
9B	PQZ	2.4 to 3.2 fl oz/A	pyrifluquinazon	1	12	L
9D	Sefina	14.0 fl oz/A	afidopyropen	0	12	L
16	Courier SC	9.0 to 13.6 fl oz/A	buprofezin	1	12	L
21A	Portal XLO	2.0 pt /A	fenpyroximate	1	12	L
23	Movento	4.0 to 5.0 fl oz/A	spirotetramat	1	24	L
23	Oberon 2SC	7.0 to 8.5 fl oz/A	spiromesifen	1	12	M
28 + 6	Minecto Pro	10.0 fl oz/A	cyantraniliprole + abamectin*	7	12	H
n/a	Requiem EC	2.0 to 3.0 qt/A	<i>Chenopodium</i> extract	0	4	L

## Group 3A Pyrethroid Insecticides Registered for Use on Tomatoes

Apply one of the following formulations (check if the product label lists the insect you intend to spray; the label is the law):						
Product Name	Product Rate	Active Ingredient(s) (* = Restricted Use)	PHI (d)	REI (h)	Bee TR	
Asana XL	2.9 to 9.6 fl oz/A	esfenvalerate*	1	12	H	
Baythroid XL <sup>1</sup>	2.1 to 2.8 fl oz/A	beta-cyfluthrin*	0	12	H	

Group 3A Pyrethroid Insecticides Registered for Use on Tomatoes Group - continued on next page



## F Tomatoes

### 3A Pyrethroid Insecticides Registered for Use on Tomatoes - continued

Brigade 2EC, others	2.1 to 5.2 fl oz/A	bifenthrin*	1	12	H
Danitol 2.4EC	10.67 fl oz/A	fenpropathrin*	3	24	H
Hero EC	4.0 to 10.3 fl oz/A	zeta-cypermethrin* + bifenthrin*	1	12	H
Lambda-Cy IEC, others	1.28 to 3.84 fl oz/A	lambda-cyhalothrin*	5	24	H
Mustang Maxx	2.24 to 4.0 fl oz/A	zeta-cypermethrin*	1	12	H
Proaxis <sup>1</sup>	1.92 to 3.84 fl oz/A	gamma-cyhalothrin*	5	24	H
Tombstone, others	1.6 to 2.8 fl oz/A	cyfluthrin*	0	12	H
Warrior II <sup>1</sup>	0.96 to 1.92 fl oz/A	lambda-cyhalothrin*	5	24	H
<b>Combo products containing a pyrethroid</b>					
Besiege	5.0 to 9.0 fl oz/A	lambda-cyhalothrin* + chlorantraniliprole (Group 28)	5	24	H
Brigadier	3.8 to 9.85 fl oz/A	bifenthrin* + imidacloprid (Group 4A) - foliar	1	12	H
Endigo ZC <sup>1</sup>	4.0 to 4.5 fl oz/A	lambda-cyhalothrin* + thiamethoxam (Group 4A)	5	24	H
Gladiator	19.0 fl oz/A	zeta-cypermethrin* + abamectin* (Group 6)	7	12	H
Leverage 360 <sup>1</sup>	3.8 to 4.1 fl oz/A	beta-cyfluthrin* + imidacloprid (Group 4A)	0	12	H
Swagger	7.6 to 19.7 fl oz/A	bifenthrin* + imidacloprid (Group 4A) - foliar	7	12	H

<sup>1</sup>Resistance concerns with Western flower thrips

### Group 4A Neonicotinoid Insecticides Registered for Use on Tomatoes

Apply one of the following formulations (check if the product label lists the insect you intend to spray; the label is the law):

Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
Admire Pro	7.0 to 10.5 fl oz/A	imidacloprid - soil	21	12	H
Admire Pro	1.3 to 2.2 fl oz/A	imidacloprid - foliar	0	12	H
Assail 30SG	1.5 to 4.0 oz/A	acetamiprid	7	12	M
Actara 25WDG	2.0 to 5.5 oz/A	thiamethoxam	0	12	H
Platinum 75SG	1.66 to 3.67 oz/A	thiamethoxam	30	12	H
Scorpion 35SL	9.0 to 10.5 fl oz/A	dinotefuran - soil	21	12	H
Scorpion 35SL	2.0 to 7.0 fl oz/A	dinotefuran - foliar	1	12	H
Venom 70SG	5.0 to 7.5 oz/A	dinotefuran - soil	21	12	H
Venom 70SG	1.0 to 4.0 oz/A	dinotefuran - foliar	1	12	H
<b>Combo products containing a neonicotinoid</b>					
Brigadier	3.8 to 9.85 fl oz/A	imidacloprid + bifenthrin* (Group 3A) - foliar	1	12	H
Durivo	10.0 to 13.0 fl oz/A	thiamethoxam + chlorantraniliprole (Group 28)	30	12	H
Endigo ZC <sup>1</sup>	4.0 to 4.5 fl oz/A	thiamethoxam + lambda-cyhalothrin* (Group 3A)	5	24	H
Leverage 360 <sup>1</sup>	3.8 to 4.1 fl oz/A	imidacloprid + beta-cyfluthrin* (Group 3A)	7	12	H
Swagger	7.6 to 19.7 fl oz/A	imidacloprid + bifenthrin* (Group 3A) - foliar	1	12	H
Voliam Flexi	4.0 to 7.0 oz/A	thiamethoxam + chlorantraniliprole	1	12	H

<sup>1</sup>Resistance concerns with Western flower thrips

## Disease Control

### THE LABEL IS THE LAW-see the Pesticide Use Disclaimer on the first page of chapter F. Recommended Fungicides

**Nematodes** See sections E 1.5 Soil Fumigation and E 1.6 Nematode Control in chapter E Pest Management.

### Seed Treatment

Purchase hot water treated seed or request hot water treatment. Heat treatment is a non-chemical alternative to conventional chlorine treatments that only kills pathogens on the surface of the seed coat. Heat treatment has the additional benefit of killing pathogens within the seed coat and is particularly useful for crops that are prone to seed-borne bacterial infections such as tomato and pepper. Seed heat treatment follows a strict time and temperature protocol and is best done with thermostatically controlled water baths. Two baths are required: one for pre-heating, and a second for the effective (pathogen killing) temperature. Seeds stay in the first bath at 100°F (38°C) for 10 minutes, and in the second bath at 122°F (50°C) for 25 minutes. Immediately after removal from the second bath, seeds should be thoroughly rinsed with cool water, and dried on a screen or paper.

Alternatively, soak seeds in a mixture of 1part Alcide (sodium chlorite), 1 part lactic acid, and 18 parts water

for 1-2 minutes under constant agitation, and rinse for 5 minutes in cool running water. Do not use pelleted seeds because moisture results in the loss of coating material.

Only treat seed that will be used during the current production season. Following heat or chlorine treatment, dust dried seed with Captan 50WP or Thiram 480DP at 1 level tsp/lb seed (3.0 oz/100 lb).

## Damping-off and Root Rots

*Greenhouse:* Use seed treatment and plant in a disease-free mix.

*Field:* At planting apply one of the fungicides via drip or banded spray. Additional field applications may be made as needed, see label for specific instructions.

Code	Product Name	Product Rate	Active Ingredient(s) (*= <b>Restricted Use</b> )	PHI (d)	REI (h)	Bee TR
4	MetaStar 2E AG	2.0-4.0 pt/A	metalaxyl	AP	48	N
4	Ridomil Gold 4SL <sup>1</sup>	1.0-2.0 pt/A <sup>1</sup>	mefenoxam	AP	48	N
4	Ultra Flourish 2E <sup>1</sup>	2.0-4.0 pt/A <sup>1</sup>	mefenoxam	AP	48	N
P07	Aliette 80WDG	2.5 to 5.0 lb/A	fosetyl-Al	14	12	N

<sup>1</sup>Apply in a 7-inch band at transplanting. Determine the amount of Ridomil Gold or Ultra Flourish per acre using the calibration formula for changing from broadcast to band application (see the section E 1.3 Calibrating Granular Applicators in chapter E Pest Management).

## Bacterial Diseases:

### Bacterial Canker

Use certified transplants. Rotate to allow 3 years between plantings. When producing transplants, use Clorox or heat-treated seed and treat used flats with sodium hypochlorite (bleach) (see section A 5 Transplant Production in the General Production Recommendations chapter). Stakes from bacterial canker infested fields should be power washed, soaked in a 20% (1part bleach plus 4 parts water) commercial bleach solution for at least 30 minutes, and power wash a second time prior to use. Avoid pruning and stringing when foliage is wet as this will promote the spread of the disease in infested fields. Applications of Actigard 50WG (0.33 oz/A increasing to 0.75 oz/A when plants are full size, see label) PLUS fixed copper (1.5 lb active/A) have been shown to reduce bacterial canker symptoms on fruit.

### Bacterial Speck and Bacterial Spot

When producing transplants, use Clorox or heat-treated seed as described above under Seed Treatment. Apply streptomycin sprays (Agri-Mycin 17, Agri-Strep, 1.0 lb/100gal, 1.25 tsp/gal) when the first true leaves appear and continue every 45 days until transplanting. Streptomycin cannot be used after transplanting. Limit handling of plants and keep greenhouse moisture levels low.

Rotate to allow 2 - 3 years between plantings. Use only certified transplants. Bacterial speck and/or spot occur more often on southern-produced transplants. Strains of copper resistant bacterial spot are common in some areas of the mid-Atlantic particularly on the Eastern Shore of VA. Use Actigard alone or in conjunction with copper-containing materials. Where disease is present or anticipated, do not work in fields when plant surfaces are wet.

Code	Product Name	Product Rate	Active Ingredient(s) (*= <b>Restricted Use</b> )	PHI (d)	REI (h)	Bee TR
<b>Tank mix the following beginning shortly after transplanting and repeat every 7 days.</b>						
M01	copper (OMRI)	1.0 lb ai/A	copper	0	see label	N
M03	mancozeb 75DF	1.5 lb/A	mancozeb	5	12/24	N
<b>And rotate with or apply the following:</b>						
M01+M03	ManKocide 61WP	2.5 to 5.0 lb/A	copper hydroxide + mancozeb	5	48	N
<b>The following is a plant defense activator and preventative applications should begin prior to the onset of symptoms.</b>						
P01	Actigard 50WG <sup>1</sup>	0.33 to 0.75 oz/A (see label)	acibenzolar-S-methyl	14	12	N

<sup>1</sup>Use in areas where copper resistance is known. See label for rates and times of use.

### Bacterial Wilt

Use certified transplants. Avoid growing tomatoes in fields where bacterial wilt has occurred. Crop rotation to non--host crops is the best measure to reduce levels of bacterial wilt. In particular, avoid planting where tomatoes or peppers were grown in the preceding year. Some resistant cultivars, such as BHN669, are available. Avoid irrigating with pond water when possible, especially for ponds that are adjacent to previously diseased fields as they may be contaminated with the causal agent.

**Fungal Diseases:****Botrytis Fruit Rot (Gray Mold)**

Gray mold is a problem during the fall in fields with dense foliage and poor drainage. For fall production, select fields with good drainage.

Code	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
<b>Shortly before harvest or when conditions are wet and cool, rotate the following as long as weather conditions favor disease development:</b>						
M05	chlorothalonil 6F	2.0 to 2.75 pt/A also very good for late blight	chlorothalonil	0	12	N
7	Endura 70W	9.0 to 12.5 oz/A also very good for early blight	boscalid	0	12	--
9 + 12	Switch 62.5WG	11.0 to 14.0 oz/A	cyprodinil + fludioxonil	0	12	L

**Buckeye Rot caused by *Phytophthora parasitica* and Fruit Rot caused by *Pythium* spp.**

Code	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
<b>Apply one of the following as a soil surface application under the vines 48 weeks before harvest. Apply broadcast or banded (adjust amount). Irrigate after application.</b>						
4	Ridomil Gold 4SL	1.0 pt/A	mefenoxam	AP	48	N
4	Ultra Flourish 2E	1.0 qt/A	mefenoxam	AP	48	N
<b>An alternative to soil application of mefenoxam: Apply one of the following as a foliar spray beginning when crown fruit are one-third their final size. repeat every 14 days up to a total of 3 times:</b>						
4 + M01	Ridomil Gold Copper 65WP	2.0 lb/A	mefenoxam + copper	14	48	--
4 + M05	Flouronil 76WP	2.0 lb/A	mefenoxam + chlorothalonil	14	48	N
4 + M05	Ridomil Gold Bravo 76WP	2.0 lb/A	mefenoxam + chlorothalonil	14	48	N
<b>If weather and soil conditions continue to favor disease development apply one of the following between applications of the above listed fungicides:</b>						
11 + 27	Tanos 50DF	8.0 oz/A	famoxadone + cymoxanil	3	12	--
22+M03	Gavel 75DF	1.5 to 2.0 lb/A	zoxamide + mancozeb	5	48	--

**Fusarium Wilt and Verticillium Wilt**

Select varieties with resistance to Fusarium and Verticillium wilts. For Fusarium wilt, select cultivars that are resistant to Races 1, 2, and 3 as all are prevalent on in the Mid-Atlantic region. Soil fumigation and proper crop rotation are essential components of a successful management program.

**Late Blight**

Use disease free transplants. If possible, produce your own transplants since transplants obtained from other regions may increase the risk of a late blight infestation. A strong scouting program, preventative fungicide applications when warranted, and microclimate management to reduce levels of free moisture on foliage are essential to help reduce the potential for disease development. Tomato cultivars with resistance to Late blight are available.

Code	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
<b>When plants are 6 inches tall, apply one of the following protectant fungicides and repeat every 7 days.</b>						
M03	mancozeb 75DF	3.0 lb/A	mancozeb	5	12,24	N
M03+22	Gavel 75DF	1.5 to 2.0 lb/A	mancozeb + zoxamide	5	48	--
M05	chlorothalonil 6F	1.0 to 3.0 pt/A	chlorothalonil	0	12	N
M05+22	Zing! 4.9SC	36.0 fl oz/A	chlorothalonil + zoxamide	7	12	N
<b>Protectant fungicides should only be applied preventatively. Monitor the movement of the disease at <a href="http://www.usablight.org/">http://www.usablight.org/</a> or via local online Extension resources. Once late blight is detected in your area, TANK MIX one of the following translaminar fungicides which can move into and through leaves WITH A PROTECTANT FUNGICIDE such as chlorothalonil, Gavel, or mancozeb. Products containing mefenoxam should not be used unless your extension professional or the aforementioned website are certain that current strains are sensitive. To achieve the best control rotate between one of the following options:</b>						
3 + 40	Revus Top 4.16SC	5.5 to 7.0 fl oz/A; also offers protection from leaf spots; <b>not</b> for use on small fruited varieties.	difenoconazole + mandipropamid	1	12	M
49+M05	Orondis Opti 3.37SC	1.75 to 2.5 pt/A; also offers protection from leaf spots	oxathiapiprolin + chlorothalonil	0	12	--
49+40	Orondis Ultra 2.33SC	5.5 to 8.0 fl oz/A	oxathiapiprolin + mandipropamid	1	4	--

Late Blight - continued on next page

## Late Blight - continued

11	Reason 500SC	5.5 to 8.2 fl oz/A	fenamidone	14	12	--
11+27	Tanos 50DF	8.0 oz/A; also offers protection from leaf spots	famoxadone + cymoxanil	3	12	--
21	Ranman 400SC	2.10 to 2.75 fl oz/A	cyazofamid	0	12	L
27	Curzate 60DF	3.2 to 5.0 oz/A	cymoxanil	3	12	N
28	Previcur Flex 6F	1.5 pt/A	propamocarb HCl	5	12	N
40	Forum 4.17SC	6.0 fl oz/A	dimethomorph	4	12	N
43	Presidio 4SC	3.0 to 4.0 fl oz/A	fluopicolide	2	12	L
<b>GREENHOUSE USE: Consult fungicide labels to ensure greenhouse applications are permitted. The following materials permit greenhouse applications and can offer suppression. Apply one of the following:</b>						
M05+P07	Catamaran 5.3F	5.5 to 7.0 pt/A	chlorothalonil + phosphite	0	12	--
11	Heritage 50WG	1.6 to 2.0 oz/A	azoxystrobin	0	4	N

## Leaf Mold

Leaf mold is caused by the fungus *Passalora fulva* (previously called *Fulvia fulva* or *Cladosporium fulvum*). Leaf mold may occur during periods of high moisture particularly within the canopy. Leaf mold is primarily damaging in greenhouse and high tunnel tomato settings with long periods of high relative humidity. Vent structures regularly to reduce humidity and leaf wetness. See Table E-11 for fungicides labeled for use in greenhouses.

Code	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
<b>Apply or rotate between the following fungicides as long as conditions are favorable for disease development:</b>						
M05+P07	Catamaran 5.3F	4.5 to 7.0 pt/A	chlorothalonil + phosphite	4	12	--
3 + 40	Revus Top 4.16SC	5.5 to 7.0 fl oz/A	difenoconazole + mandipropamid	1	12	M

## Leaf Spots caused by Early blight and Septoria leaf spot and Fruit Rots caused by Anthracnose and Early blight:

Follow a crop rotation with at least 2 years without tomatoes or potatoes. Use disease-free transplants and disease resistant varieties when possible. In high elevated areas, in fields not rotated away from tomatoes, or in late planted fields begin sprays shortly after transplanting. In all other areas, follow a regular (7-day) spray schedule.

Code	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
<b>Alternate or tank mix one of the following protectant fungicides:</b>						
M05+49	Orondis Opti 3.37SC	1.75 to 2.5 pt/A (also for late blight)	chlorothalonil + oxathiapiprolin	0	12	--
M03	mancozeb 75DF	3.0 lb/A (also for gray leaf spot and leaf mold)	mancozeb	5	12/24	N
M03+22	Gavel 75DF	1.5 to 2.0 lb/A	mancozeb + zoxamide	5	48	--
M05	chlorothalonil 6F	2.0 to 3.0 pt/A (also for gray leaf spot, black mold and soil rot)	chlorothalonil	0	12	N
M05+22	Zing! 4.9SC	36.0 fl oz/A	chlorothalonil + zoxamide	7	12	N
<b>WITH one of the following fungicides (fungicides from different FRAC codes should be rotated to help reduce the chances for fungicide resistance development):</b>						
3 + 7	Aprovia Top 1.62EC	10.5 to 13.5 fl oz/A	difenoconazole + benzovindiflupyr	0	12	--
3 + 9	Inspire Super 2.82EW	16.0 to 20.0 fl oz/A	difenoconazole + cyprodonil	0	12	--
3 + 11	Quadris Top 1.67SC	8.0 fl oz/A	difenoconazole + azoxystrobin	0	12	--
3 + 40	Revus Top 4.16SC	5.5 to 7.0 fl oz/A	difenoconazole + mandipropamid	1	12	M
7	Endura 70W	2.5 to 3.5 oz/A (also for <i>Botrytis</i> at 9.0 to 12.5 oz/A)	boscalid	3	12	--
3 + 11	Topguard 4.29SC	4.0 to 8.0 fl oz/A	flutriafol + azoxystrobin	0	12	--
7	Fontelis 1.67SC	16.0 to 24.0 fl oz/A	penthiopyrad	0	12	L
7 + 11	Priaxor 4.17SC	4.0 to 8.0 fl oz/A	fluxapyroxad + pyraclostrobin	7	12	N
11	azoxystrobin 2.08F	5.0 to 6.2 fl oz/A (also for black mold and buckeye rot)	azoxystrobin ( <b>Do not apply near apples</b> , see label)	0	4	N
11	Cabrio 20EG	8.0 to 12.0 oz/A	pyraclostrobin	0	12	N
11	Flint Extra 500SC	3.0-3.8 fl oz/A	trifloxystrobin ( <b>Do not apply near Concord grapes</b> , see label)	3	12	N
11 + 27	Tanos 50DF	8.0 oz/A PLUS protectant fungicide (also for buckeye rot suppression and gray leaf spot).	famoxadone + cymoxanil	12	3	--

F Tomatoes

**Postharvest Rots**

Avoid harvesting when the foliage is wet. To prevent rots in mature green tomatoes, avoid washing freshly harvested fruit in cold water. To prevent movement of bacteria into the stem end of the fruit, do not allow water temperatures in flumes and tanks of more than 10°F above fruit temperature. Use a minimum of 100 ppm free chlorine and keep pH between 6.5-7.0 in the flume. Store at 55°F (13°C) with relative humidity of 80%. For more information on postharvest tomato diseases, see <http://edis.ifas.ufl.edu/HS131>.

**Powdery Mildew**

Code	Product Name	Product Rate	Active Ingredient(s) (*= <b>Restricted Use</b> )	PHI (d)	REI (h)	Bee TR
<b>At first appearance of the disease, rotate between the following fungicides<sup>1</sup>:</b>						
<b>FIELD, repeat every 7 to 14 days:</b>						
3	Rally 40WSP	2.5 to 4.0 oz/A	myclobutanil	0	12	N
3 + 7	Aprovia Top 1.62EC	10.5 to 13.5 fl oz/A	difenoconazole + benzovindiflupyr	0	12	--
3 + 9	Inspire Super 2.82EW	16.0 to 20.0 fl oz/A	difenoconazole + cyprodonil	0	12	--
3 + 40	Revus Top 4.16SC	5.5 to 7.0 fl oz/A	difenoconazole + mandipropamid	1	12	M
11	Cabrio 20EG	8.0 to 12.0 oz/A	pyraclostrobin	0	12	N
13	Quintec 2.08SC	6.0 fl oz/A	quinoxifen	3	12	--
<b>GREENHOUSE<sup>2</sup>, thoroughly cover upper and lower leaf surfaces and repeat every 7 days:</b>						
--	JMS Stylet-Oil	1.0 to 2.0 gal/100 gal	paraffinic oil	--	--	--
9	Scala 5SC	7.0 fl oz/A	pyrimethanil	1	12	--

<sup>1</sup>Fungicides from different FRAC codes should be rotated to help reduce the chances for fungicide resistance development. <sup>2</sup>Powdery mildew can cause serious problems in greenhouse and high tunnel settings. See Table E-14 for additional fungicides labeled for use in greenhouses.

**Southern Blight (*Sclerotium rolfsii*)**

Southern blight is most commonly seen in the southern part of the Mid-Atlantic region. High soil moisture and temperatures favor disease, while long crop rotations with corn and small grains help reduce disease incidence. Weed control is important as *Sclerotium rolfsii* can infect a number of common weeds in the Mid-Atlantic region. Soil fumigation and staking will greatly reduce disease incidence. Applications of Blocker 4F in transplant water or as an in-furrow treatment may suppress the disease.

Code	Product Name	Product Rate	Active Ingredient(s) (*= <b>Restricted Use</b> )	PHI (d)	REI (h)	Bee TR
14	Blocker 4F	See label	pentachloronitrobenzene (PCNB)	AP	12	H

**Timber Rot (*Sclerotinia sclerotiorum*)**

Tomato timber rot, also known as sclerotinia stem rot, is a fungal disease caused by *Sclerotinia sclerotiorum*. Rotate away from fields where snap or lima beans, peas, peanuts, lettuce, or cucurbits were grown in the past. -- Timber rot occurs during prolonged wet periods and cooler temperatures (<80°F).

**Viruses: Tomato Spotted Wilt Virus (TSWV)**

TSWV can result in severely stunted plants. The virus is spread by thrips from ornamental flowering plants, field crops, and weeds to tomatoes. TSWV can be particularly devastating in southern and eastern parts of VA. Use resistant varieties when available. Do not grow any ornamental bedding plants in the same greenhouse as tomato transplants. Control weeds in and around greenhouses, high tunnels, or transplant areas. Monitor greenhouses and tomato fields for thrips and begin an insecticide control program once observed. Use of reflective mulch can help repel thrips. If tomato crops are near wheat or barley fields be aware of increased thrips pressure once these crops start to turn brown in the spring.

**For Immediate Medical Attention**

**Call 911**

**For a Pesticide Exposure Poisoning  
Emergency Call**



**For All States**

This number will automatically connect you to the poison center nearest to you.

**Anyone with a poisoning emergency can call the toll-free telephone number for help.** Personnel at the Center will give you first-aid information and direct you to local treatment centers if necessary.

## **For Pesticide Spills**

**Small Spills:** See the product label for cleanup advice.

**Large spills:** Call the National Response Center at 1-800-424-8802 or CHEMTREC at 800-424-9300 (24 hours) - Industry assistance with emergency response cleanup procedures for large, dangerous spills.

**Be aware of your responsibility to report spills to the proper state agency.**